

Essence: Bonn reduced.

Homomorphic Expansions and w-Knots

Dror Bar-Natan, UWO February 2010, <http://www.math.toronto.edu/~drorbn/Talks/UWO-100225/>

"God created the knots, all else in topology is the work of mortals."
Leopold Kronecker (modified)

What are w-Trivalent Tangles? (PA := Planar Algebra)

$\{\text{knots} \& \text{links}\} = \text{PA} \left\langle \begin{array}{l} \text{R123} : \text{ } \end{array} \middle| \text{ } \right\rangle_{0 \text{ legs}}$

$\{\text{trivalent tangles}\} = \text{PA} \left\langle \begin{array}{l} \text{R23, R4} : \end{array} \middle| \text{ } \right\rangle$

wTT =

$\{\text{trivalent w-tangles}\} = \text{PA} \left\langle \begin{array}{l} \text{w-} \\ \text{generators} \end{array} \middle| \begin{array}{l} \text{w-} \\ \text{relations} \end{array} \middle| \begin{array}{l} \text{unary w-} \\ \text{operations} \end{array} \right\rangle$

The w-generators.

Homomorphic expansions for a filtered algebraic structure \mathcal{K} :

$$\text{ops} \supset \mathcal{K} = \mathcal{K}_0 \supset \mathcal{K}_1 \supset \mathcal{K}_2 \supset \mathcal{K}_3 \supset \dots$$

$$\downarrow \qquad \qquad \qquad \downarrow Z$$

$$\text{ops} \supset \text{gr } \mathcal{K} := \mathcal{K}_0/\mathcal{K}_1 \oplus \mathcal{K}_1/\mathcal{K}_2 \oplus \mathcal{K}_2/\mathcal{K}_3 \oplus \mathcal{K}_3/\mathcal{K}_4 \oplus \dots$$

An **expansion** is a filtration respecting $Z : \mathcal{K} \rightarrow \text{gr } \mathcal{K}$ that "covers" the identity on $\text{gr } \mathcal{K}$. A **homomorphic expansion** is an expansion that respects all relevant "extra" operations.

Filtered algebraic structures are cheap and plenty. In any \mathcal{K} , allow formal linear combinations, let \mathcal{K}_1 be the ideal generated by differences (the "augmentation ideal"), and let $\mathcal{K}_m := \langle (\mathcal{K}_1)^m \rangle$ (using all available "products").

"An Algebraic Structure"

- Has kinds, objects, operations, and maybe constants.
- Perhaps subject to some axioms.
- We always allow formal linear combinations.

Example: Pure Braids. PB_n is generated by x_{ij} , "strand i goes around strand j once", modulo "Reidemeister moves". $A_n := \text{gr } PB_n$ is generated by $t_{ij} := x_{ij} - 1$, modulo the $4T$ relations $[t_{ij}, t_{ik} + t_{jk}] = 0$ (and some lesser ones too). Much happens in A_n , including the Drinfel'd theory of associators.

Our case(s).

$$\mathcal{K} \xrightarrow[\text{solving finitely many equations in finitely many unknowns}]{Z: \text{high algebra}} \mathcal{A} := \text{gr } \mathcal{K} \xrightarrow[\text{low algebra: pictures represent formulas}]{\text{given a "Lie" algebra } \mathfrak{g}} \mathcal{U}(\mathfrak{g})$$

\mathcal{K} is knot theory or topology; $\text{gr } \mathcal{K}$ is finite combinatorics; bounded-complexity diagrams modulo simple relations.

[1] <http://qlink.queensu.ca/~4lb11/interesting.html> 8/2/10, 10:02pm
Also see <http://www.math.toronto.edu/~drorbn/papers/WKO/>

A Ribbon 2-Knot is a surface S embedded in \mathbb{R}^4 that bounds an immersed handlebody B , with only "ribbon singularities"; a ribbon singularity is a disk D of transverse double points, whose preimages in B are a disk D_1 in the interior of B and a disk D_2 with $D_2 \cap \partial B = \partial D_2$, modulo isotopies of S alone.

The w-relations include R234, VR1234, M, Overcrossings Commute (OC) but not UC, $W^2 = 1$, and funny interactions between the wen and the cap and over- and under-crossings:

Challenge. Do the Reidemeister!

The unary w-operations

Just for fun.

An expansion Z is a choice of a "progressive scan" algorithm.

$$\mathcal{K}/\mathcal{K}_1 \oplus \mathcal{K}_1/\mathcal{K}_2 \oplus \mathcal{K}_2/\mathcal{K}_3 \oplus \mathcal{K}_3/\mathcal{K}_4 \oplus \mathcal{K}_4/\mathcal{K}_5 \oplus \mathcal{K}_5/\mathcal{K}_6 \oplus \dots$$

$$\parallel \qquad \qquad \qquad \parallel$$


$$\mathbb{R} \qquad \qquad \qquad \ker(\mathcal{K}/\mathcal{K}_4 \rightarrow \mathcal{K}/\mathcal{K}_3)$$

Crop Rotate Adjoin


2010-02 Page 1

Homomorphic Expansions and w-Knots, Page 2

Knot-Theoretic statement. There exists a homomorphic expansion Z for trivalent w-tangles. In particular, Z should respect $R4$ and intertwine annulus and disk unzips:



Diagrammatic statement. Let $R = \exp \mathbb{H} \in \mathcal{A}^w(\uparrow\uparrow)$. There exist $\omega \in \mathcal{A}^w(\uparrow)$ and $V \in \mathcal{A}^w(\uparrow\uparrow)$ so that



Algebraic statement. With $I\mathfrak{g} := \mathfrak{g}^* \times \mathfrak{g}$, with $c : \mathcal{U}(I\mathfrak{g}) \rightarrow \hat{\mathcal{U}}(I\mathfrak{g})/\hat{\mathcal{U}}(\mathfrak{g}) = \hat{S}(\mathfrak{g}^*)$ the obvious projection, with S the antipode of $\hat{\mathcal{U}}(I\mathfrak{g})$, with W the automorphism of $\hat{\mathcal{U}}(I\mathfrak{g})$ induced by flipping the sign of \mathfrak{g}^* , with $r \in \mathfrak{g}^* \otimes \mathfrak{g}$ the identity element and with $R = e^r \in \hat{\mathcal{U}}(I\mathfrak{g}) \otimes \hat{\mathcal{U}}(\mathfrak{g})$ there exist $\omega \in \hat{S}(\mathfrak{g}^*)$ and $V \in \hat{\mathcal{U}}(I\mathfrak{g})^{\otimes 2}$ so that

(1) $V(\Delta \otimes 1)(R) = R^{13}R^{23}V$ in $\hat{\mathcal{U}}(I\mathfrak{g})^{\otimes 2} \otimes \hat{\mathcal{U}}(\mathfrak{g})$
 (2) $V \cdot SWV = 1$ (3) $(c \otimes c)(V \Delta(\omega)) = \omega \otimes \omega$

Unitary statement. There exists $\omega \in \text{Fun}(\mathfrak{g})^G$ and an (infinite order) tangential differential operator V defined on $\text{Fun}(\mathfrak{g}_x \times \mathfrak{g}_y)$ so that

(1) $V e^{x+y} = e^{\hat{x}} \hat{e}^y V$ (allowing $\hat{\mathcal{U}}(\mathfrak{g})$ -valued functions)
 (2) $V V^* = I$ (3) $V \omega_{x+y} = \omega_x \omega_y$


Group-Algebra statement. There exists $\omega^2 \in \text{Fun}(\mathfrak{g})^G$ so that for every $\phi, \psi \in \text{Fun}(\mathfrak{g})^G$ (with small support), the following holds in $\hat{\mathcal{U}}(\mathfrak{g})$:

$$\iint_{\mathfrak{g} \times \mathfrak{g}} \phi(x)\psi(y)\omega_{x+y}^2 e^{x+y} = \iint_{\mathfrak{g} \times \mathfrak{g}} \phi(x)\psi(y)\omega_x^2 \omega_y^2 e^x e^y. \quad (\text{shhh, this is Duflo})$$


Convolutions statement (Kashiwara-Vergne). Convolutions of invariant functions on a Lie group agree with convolutions of invariant functions on its Lie algebra. More accurately, let G be a finite dimensional Lie group and let \mathfrak{g} be its Lie algebra, let $j : \mathfrak{g} \rightarrow \mathbb{R}$ be the Jacobian of the exponential map $\exp : \mathfrak{g} \rightarrow G$, and let $\Phi : \text{Fun}(G) \rightarrow \text{Fun}(\mathfrak{g})$ be given by $\Phi(f)(x) := j^{1/2}(x)f(\exp x)$. Then if $f, g \in \text{Fun}(G)$ are Ad-invariant and supported near the identity, then

$$\Phi(f) \star \Phi(g) = \Phi(f \star g).$$

From $w\Gamma$ to \mathcal{A}^w . $\text{gr}_m w\Gamma := \{m\text{-cubes}\}/\{(m+1)\text{-cubes}\}$:



w-Jacobi diagrams and \mathcal{A} . $\mathcal{A}^w(Y \uparrow) \cong \mathcal{A}^w(\uparrow\uparrow)$ is



Diagrammatic to Algebraic. With (x_i) and (φ^j) dual bases of \mathfrak{g} and \mathfrak{g}^* and with $[x_i, x_j] = \sum b_{ij}^k x_k$, we have $\mathcal{A}^w \rightarrow \mathcal{U}$ via

$$\sum_{i,j,k,l,m,n=1}^{\dim \mathfrak{g}} b_{ij}^k b_{kl}^m \varphi^i \varphi^j x_n x_m \varphi^n \varphi^l \in \mathcal{U}(I\mathfrak{g})$$

Unitary \iff Algebraic. The key is to interpret $\hat{\mathcal{U}}(I\mathfrak{g})$ as tangential differential operators on $\text{Fun}(\mathfrak{g})$:

- $\varphi \in \mathfrak{g}^*$ becomes a multiplication operator.
- $x \in \mathfrak{g}$ becomes a tangential derivation, in the direction of the action of $\text{ad } x : (x\varphi)(y) := \varphi([x, y])$.
- $c : \hat{\mathcal{U}}(I\mathfrak{g}) \rightarrow \hat{\mathcal{U}}(I\mathfrak{g})/\hat{\mathcal{U}}(\mathfrak{g}) = \hat{S}(\mathfrak{g}^*)$ is "the constant term".

Unitary \implies Group-Algebra. $\iint \omega_{x+y}^2 e^{x+y} \phi(x)\psi(y)$

$$= \langle \omega_{x+y}, \omega_{x+y} e^{x+y} \phi(x)\psi(y) \rangle = \langle V \omega_{x+y}, V e^{x+y} \phi(x)\psi(y) \omega_{x+y} \rangle$$

$$= \langle \omega_x \omega_y, e^x e^y V \phi(x)\psi(y) \omega_{x+y} \rangle = \langle \omega_x \omega_y, e^x e^y \phi(x)\psi(y) \omega_x \omega_y \rangle$$

$$= \iint \omega_x^2 \omega_y^2 e^x e^y \phi(x)\psi(y).$$

Convolutions and Group Algebras (ignoring all Jacobians). If G is finite, A is an algebra, $\tau : G \rightarrow A$ is multiplicative then

$$\text{Fun}(G, \star) \cong (A, \cdot) \text{ via } L : f \mapsto \sum f(a)\tau(a).$$

$$\begin{array}{ccc} (\mathfrak{g}, +) \ni x & \xrightarrow{\tau_0 = \exp_G} & e^x \in \hat{S}(\mathfrak{g}) \\ \downarrow \exp_G & \searrow \exp_{\mathfrak{g}} & \downarrow \chi \\ (G, \cdot) \ni e^x & \xrightarrow{\tau_1} & e^x \in \hat{\mathcal{U}}(\mathfrak{g}) \end{array} \quad \text{so} \quad \begin{array}{ccc} \text{Fun}(\mathfrak{g}) & \xrightarrow{L_0} & \hat{S}(\mathfrak{g}) \\ \downarrow \Phi^{-1} & & \downarrow \chi \\ \text{Fun}(G) & \xrightarrow{L_1} & \hat{\mathcal{U}}(\mathfrak{g}) \end{array}$$

with $L_0 \psi = \int \psi(x) e^x dx \in \hat{S}(\mathfrak{g})$ and $L_1 \Phi^{-1} \psi = \int \psi(x) e^x \in \hat{\mathcal{U}}(\mathfrak{g})$. Given $\psi_1 \in \text{Fun}(\mathfrak{g})$ compare $\Phi^{-1}(\psi_1) \star \Phi^{-1}(\psi_2)$ and $\Phi^{-1}(\psi_1 \star \psi_2)$ in $\hat{\mathcal{U}}(\mathfrak{g})$: (shhh, $L_{0,1}$ are "Laplace transforms")

$$\star \text{ in } G : \iint \psi_1(x)\psi_2(y) e^x e^y \quad \star \text{ in } \mathfrak{g} : \iint \psi_1(x)\psi_2(y) e^{x+y}$$

- We skipped...**
- The Alexander polynomial and Milnor numbers.
 - u-Knots, Alekseev-Torossian, and Drinfel'd associators.
 - v-Knots, quantum groups and Etingof-Kazhdan.
 - BF theory and the successful religion of path integrals.
 - The simplest problem hyperbolic geometry solves.