

Improve PNG resolution!
consider doing topology before algebra, after all.
say something explicit about AT-KV?

Trees and Wheels and Balloons and Hoops

Dror Bar-Natan, Toronto, March 2013
 $\omega\epsilon\beta := \text{http://www.math.toronto.edu/~drorbn/Talks/Toronto-1303}$

15 Minutes on Algebra

Let T be a finite set of "tail labels" and H a finite set of "head labels". Set

$$M_{1/2}(T; H) := FL(T)^H,$$

" H -labeled lists of elements of the degree-completed free Lie algebra generated by T ".

$$FL(T) = \left\{ 2t_2 - \frac{1}{2}[t_1, [t_1, t_2]] + \dots \right\} / \left(\begin{array}{l} \text{anti-symmetry} \\ \text{Jacobi} \end{array} \right)$$

... with the obvious bracket.

$$M_{1/2}(u, v; x, y) = \left\{ \lambda = \left(x \rightarrow \begin{array}{c} u \quad v \\ \diagdown \quad \diagup \\ x \end{array}, y \rightarrow \begin{array}{c} v \quad u \\ \diagdown \quad \diagup \\ y \end{array} - \frac{22}{7} \begin{array}{c} u \quad u \quad v \\ \diagdown \quad \diagup \\ y \end{array} \right) \dots \right\}$$

Operations $M_{1/2} \rightarrow M_{1/2}$. newspeak!

Tail Multiply tm_{uv}^{uv} is $\lambda \mapsto \lambda \parallel (u, v \rightarrow w)$, satisfies "meta-associativity", $tm_{uv}^{uv} \parallel tm_{uv}^{uv} = tm_{uv}^{uv} \parallel tm_{uv}^{uv}$.

Head Multiply hm_z^{xy} is $\lambda \mapsto (\lambda \setminus \{x, y\}) \cup (z \rightarrow \text{bch}(\lambda_x, \lambda_y))$, where

$$\text{bch}(\alpha, \beta) := \log(e^\alpha e^\beta) = \alpha + \beta + \frac{[\alpha, \beta]}{2} + \frac{[\alpha, [\alpha, \beta]] + [[\alpha, \beta], \beta]}{12} + \dots$$

satisfies $\text{bch}(\text{bch}(\alpha, \beta), \gamma) = \log(e^{\text{bch}(\alpha, \beta)} e^\gamma) = \text{bch}(\alpha, \text{bch}(\beta, \gamma))$, and hence meta-associativity, $hm_x^{xy} \parallel hm_x^{xz} = hm_y^{yz} \parallel hm_x^{xy}$.

Tail by Head Action tha^{ux} is $\lambda \mapsto \lambda \parallel RC_u^{\lambda_x}$, where $C_u^{-\gamma}: FL \rightarrow FL$ is the substitution $u \rightarrow e^{-\gamma} u e^\gamma$, or more precisely,


$$C_u^{-\gamma}: u \rightarrow e^{-\text{ad} \gamma}(u) = u - [\gamma, u] + \frac{1}{2}[\gamma, [\gamma, u]] - \dots,$$

and RC_u^γ is the inverse of that. Note that $C_u^{\text{bch}(\alpha, \beta)} = C_u^\alpha \parallel RC_u^\beta \parallel C_u^\alpha$ and hence "meta $u^{xy} = (u^x)^y$ ",

$$hm_z^{xy} \parallel tha^{uz} = tha^{ux} \parallel tha^{uy} \parallel hm_z^{xy},$$

and $tm_{uv}^{uv} \parallel C_w^\gamma \parallel tm_{uv}^{uv} = C_u^\gamma \parallel RC_u^{-\gamma} \parallel C_u^\gamma \parallel tm_{uv}^{uv}$ and hence "meta $(uv)^x = u^x v^x$ ", $tm_{uv}^{uv} \parallel tha^{wx} = tha^{ux} \parallel tha^{vx} \parallel tm_{uv}^{uv}$.


Wheels. Let $M(T; H) := M_{1/2}(T; H) \times CW(T)$, where $CW(T)$ is the (completed graded) vector space of cyclic words on T , or equally well, on $FL(T)$:



Operations. On $M(T; H)$, define tm_{uv}^{uv} and hm_z^{xy} as before, and tha^{ux} by adding some J -spice:

$$(\lambda; \omega) \mapsto (\lambda, \omega + J_u(\lambda_x)) \parallel RC_u^\gamma,$$

where $J_u(\gamma) := \int_0^1 ds \text{div}_u(\gamma \parallel RC_u^{s\gamma}) \parallel C_u^{-s\gamma}$, and




Theorem Green. All green identities still hold.

Merge Operation. $(\lambda_1; \omega_1) * (\lambda_2; \omega_2) := (\lambda_1 \cup \lambda_2; \omega_1 + \omega_2)$.


15 Minutes on Topology

$\mathcal{K}^{bh}(T; H)$.

"Ribbon-knotted balloons and hoops"




Examples.



More on δ . δ maps v-tangles to \mathcal{K}^{bh} ; the kernel is as above, and conjecturally, that's all. Allowing punctures and cuts, δ is onto.

satisfies R123, VR123, D, and




δ injects u-knots into \mathcal{K}^{bh} (likely u-tangles too). δ maps v-tangles to \mathcal{K}^{bh} ; the kernel is as above, and conjecturally, that's all. Allowing punctures and cuts, δ is onto.

Operations

Punctures & Cuts. Connected Sums. $(\text{tree with balloon}) * (\text{tree with balloon}) = \text{tree with two balloons}$


If X is a space, $\pi_1(X)$ is a group, $\pi_2(X)$ is an Abelian group, and π_1 acts on π_2 .



Properties

- Associativities: $m_a^{ab} \parallel m_a^{ac} = m_b^{bc} \parallel m_a^{ab}$, for $m = tm, hm$.
- " $(uv)^x = u^x v^x$ ": $tm_{uv}^{uv} \parallel tha^{wx} = tha^{ux} \parallel tha^{vx} \parallel tm_{uv}^{uv}$,
- " $u(xy) = (u^x)^y$ ": $hm_z^{xy} \parallel tha^{uz} = tha^{ux} \parallel tha^{uy} \parallel hm_z^{xy}$.

Tangle concatenations $\rightarrow \pi_1 \times \pi_2$. With $dm_c^{ab} := tha^{ab} \parallel tm_c^{ab} \parallel hm_c^{ab}$,



Moral. To construct an M -valued invariant ζ of (v-)tangles, and nearly an invariant on \mathcal{K}^{bh} , it is enough to declare ζ on the generators, and verify the relations that δ satisfies.

Trees and Wheels and Balloons and Hoops: Why I Care

~~studies time, wko, up~~

The Invariant ζ . Set $\zeta(\epsilon_x) = (x \rightarrow 0; 0)$, $\zeta(\epsilon_u) = ((); 0)$, and

$$\zeta: \begin{array}{c} \text{circle with } u \text{ and } x \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \mapsto \begin{pmatrix} u \\ \downarrow \\ v_x \\ ; 0 \end{pmatrix} \quad \begin{array}{c} \text{circle with } x \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \mapsto \begin{pmatrix} - \\ \downarrow \\ v_x \\ ; 0 \end{pmatrix}$$

Theorem. ζ is (log of) the unique homomorphic universal finite type invariant on \mathcal{K}^{bh} .
 (... and is the tip of an iceberg)

Paper in progress with Danco, $\omega\epsilon\beta/wko$

The β quotient is M divided by all relations that universally hold when \mathfrak{g} is the 2D non-Abelian Lie algebra. Let $R = \mathbb{Q}[\{c_u\}_{u \in T}]$ and $L_\beta := R \otimes T$ with central R and with $[u, v] = c_u v - c_v u$ for $u, v \in T$. Then $FL \rightarrow L_\beta$ and $CW \rightarrow R$. Under this,

$$\mu \rightarrow ((\lambda_x); \omega) \quad \text{with } \lambda_x = \sum_{u \in T} \lambda_{ux} u x, \quad \lambda_{ux}, \omega \in R,$$

$$\text{bch}(u, v) \rightarrow \frac{c_u + c_v}{e^{c_u + c_v} - 1} \left(\frac{e^{c_u} - 1}{c_u} u + e^{c_v} \frac{e^{c_v} - 1}{c_v} v \right),$$

if $\gamma = \sum \gamma_v v$ then with $c_\gamma := \sum \gamma_v c_v$,

$u \parallel RC_\gamma^u = \left(1 + c_u \gamma_u \frac{e^{c_\gamma} - 1}{c_\gamma}\right)^{-1} \left(e^{c_\gamma} u - c_u \frac{e^{c_\gamma} - 1}{c_\gamma} \sum_{v \neq u} \gamma_v v \right)$.

$\text{div}_u \gamma = c_u \gamma_u$, and $J_u(\gamma) = \log \left(1 + \frac{e^{c_\gamma} - 1}{c_\gamma} c_u \gamma_u\right)$, so ζ is formula-computable to all orders! **Can we simplify?**

Repackaging. Given $((x \rightarrow \lambda_{ux}); \omega)$, set $c_x := \sum_v c_v \lambda_{vx}$, replace $\lambda_{ux} \rightarrow \alpha_{ux} := c_u \lambda_{ux} \frac{e^{c_x} - 1}{c_x}$ and $\omega \rightarrow e^\omega$, use $t_u = e^{c_u}$, and write α_{ux} as a matrix. Get " **β calculus**".

β Calculus. Let $\beta(H, T)$ be

$$\left\{ \begin{array}{c} \omega \mid \begin{array}{ccc} x & y & \dots \\ \alpha_{ux} & \alpha_{uy} & \dots \\ v & \alpha_{vx} & \alpha_{vy} & \dots \\ \vdots & \dots & \dots & \dots \end{array} \mid \omega \text{ and the } \alpha_{ux} \text{'s are} \\ \text{rational functions in} \\ \text{variables } t_u, \text{ one for} \\ \text{each } u \in T. \end{array} \right\}$$

$$tm_{wv}^{uw} : \begin{array}{c} \omega \mid \dots \\ u \mid \alpha \\ v \mid \beta \\ \vdots \mid \gamma \end{array} \mapsto \begin{array}{c} \omega \mid \dots \\ \alpha + \beta \\ \gamma \end{array}, \quad \begin{array}{c} \omega_1 \mid H_1 \mid \omega_2 \mid H_2 \\ T_1 \mid \alpha_1 \mid T_2 \mid \alpha_2 \\ \omega_1 \omega_2 \mid H_1 \mid H_2 \\ T_1 \mid \alpha_1 \mid 0 \\ T_2 \mid 0 \mid \alpha_2 \end{array}$$

$$hm_z^{xy} : \begin{array}{c} \omega \mid x \ y \ \dots \\ \alpha \ \beta \ \gamma \end{array} \mapsto \begin{array}{c} \omega \mid z \ \dots \\ \alpha + \beta + \langle \alpha \rangle \beta \ \gamma \end{array}$$

$$tha^{ux} : \begin{array}{c} \omega \mid x \ \dots \\ u \mid \alpha \ \beta \\ \vdots \mid \gamma \ \delta \end{array} \mapsto \begin{array}{c} \omega \epsilon \mid x \ \dots \\ \alpha(1 + \langle \gamma \rangle / \epsilon) \ \beta(1 + \langle \gamma \rangle / \epsilon) \\ \gamma / \epsilon \ \delta - \gamma \beta / \epsilon \end{array}$$

where $\epsilon := 1 + \alpha$, $\langle \alpha \rangle := \sum_v \alpha_v$, and $\langle \gamma \rangle := \sum_{v \neq u} \gamma_v$, and let


$$R_{ux}^+ := \frac{1}{u} \mid \begin{array}{c} x \\ t_u - 1 \end{array} \quad R_{ux}^- := \frac{1}{u} \mid \begin{array}{c} x \\ t_u^{-1} - 1 \end{array}$$

On long knots, ω is the Alexander polynomial!

Why happy? An ultimate Alexander invariant: Manifestly polynomial (time and size) extension of the (multivariable) Alexander polynomial to tangles. Every step of the computation is the computation of the invariant of some topological thing (no fishy Gaussian elimination!). *If there should be an Alexander invariant with an algebraic categorification, it is this one!*

See also $\omega\epsilon\beta/regina$, $\omega\epsilon\beta/gua$. **May class:** $\omega\epsilon\beta/aarhus$

Paper in progress: $\omega\epsilon\beta/kbh$ **Class next year:** $\omega\epsilon\beta/1350$


Leopold Kronecker (modified) "God created the knots, all else in topology is the work of mortals."
www.katlas.org 

Instr/graphic version


with summi;

Calc, nit/ton

wk: ten wiss, bonn, portfolio



ζ is computable! ζ of the Borromean tangle, to degree 5:



Tensorial Interpretation. Let \mathfrak{g} be a finite dimensional Lie algebra (any!). Then there's $\tau : FL(T) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \mathfrak{g})$ and $\tau : CW(T) \rightarrow \text{Fun}(\oplus_T \mathfrak{g})$. Together, $\tau : M(T; H) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \oplus_H \mathfrak{g})$, and hence

$$e^\tau : M(T; H) \rightarrow \text{Fun}(\oplus_T \mathfrak{g} \rightarrow \mathcal{U}^{\otimes H}(\mathfrak{g})).$$

ζ and BF Theory. (See Cattaneo-Rossi, arXiv:math-ph/0210037) Let A denote a \mathfrak{g} -connection on S^4 with curvature F_A , and B a \mathfrak{g}^* -valued 2-form on S^4 . For a hoop γ_x , let $\text{hol}_{\gamma_x}(A) \in \mathcal{U}(\mathfrak{g})$ be the holonomy of A along γ_x . For a ball γ_u , let $\mathcal{O}_{\gamma_u}(B) \in \mathfrak{g}^*$ be (roughly) the integral of B (transported via A to ∞) on γ_u .

Loose Conjecture. For $\gamma \in \mathcal{K}(T; H)$,

$$\int \mathcal{D}A \mathcal{D}B e^{\int B \wedge F_A} \prod_u e^{\mathcal{O}_{\gamma_u}(B)} \bigotimes_x \text{hol}_{\gamma_x}(A) = e^\tau(\zeta(\gamma)).$$

That is, ζ is a complete evaluation of the BF TQFT.



Cattaneo



Calc, nit/ton