




## Some very good formulas for the Alexander polynomial, 2


### Operations

Punctures & Cuts



If  $X$  is a space,  $\pi_1(X)$  is a group,  $\pi_2(X)$  is an Abelian group, and  $\pi_1$  acts on  $\pi_2$ .

$K \parallel tm_w^{xy}$ :



**Definition.**  $l_{xu}$  is the linking number of hoop  $x$  with balloon  $u$ . For  $x \in H$ ,  $\sigma_x := \prod_{u \in T} T_u^{l_{xu}} \in R = R_T = \mathbb{Z}(T_a)_{a \in T}$ , the ring of rational functions in  $T$  variables.

**Theorem 2 [BNS].**  $\exists!$  an invariant  $\beta: \{\text{w-balloon and hoops}\} \rightarrow R \times M_{T \times H}(R)$ , intertwining

$$1. \left( \begin{array}{c|cc} \omega_1 & H_1 \\ \hline T_1 & A_1 \end{array}, \begin{array}{c|cc} \omega_2 & H_2 \\ \hline T_2 & A_2 \end{array} \right) \xrightarrow{\sqcup} \begin{array}{c|cc} \omega_1 \omega_2 & H_1 & H_2 \\ \hline T_1 & A_1 & 0 \\ T_2 & 0 & A_2 \end{array},$$

$$2. \frac{\omega}{\begin{matrix} u \\ v \\ T \end{matrix}} \xrightarrow{tm_w^{xy}} \frac{\omega}{\begin{matrix} w \\ w \\ T \end{matrix}} \xrightarrow{T_u, T_v \rightarrow T_w},$$

$$3. \frac{\omega}{\begin{matrix} T \\ \alpha \\ \beta \\ \Xi \end{matrix}} \xrightarrow{hm_z^{xy}} \frac{\omega}{\begin{matrix} T \\ \alpha + \sigma_x \beta \\ \Xi \end{matrix}},$$

$$4. \frac{\omega}{\begin{matrix} u \\ \alpha \\ \theta \\ T \\ \phi \\ \Xi \end{matrix}} \xrightarrow{tha^{ux}} \frac{\nu \omega}{\begin{matrix} u \\ \sigma_x \alpha / \nu \\ \sigma_x \theta / \nu \\ T \\ \phi / \nu \\ \Xi - \phi / \nu \end{matrix}},$$

and satisfying  $(\epsilon_x, \epsilon_u, \rho_{ux}^\pm) \xrightarrow{\beta} \left( \frac{1}{u} \left| \begin{matrix} x \\ u \end{matrix} \right. ; \frac{1}{u} \left| \begin{matrix} x \\ T_u^{\pm 1} - 1 \end{matrix} \right. \right)$ .



**Proposition.** If  $T$  is a u-tangle and  $\beta(\delta T) = (\omega, A)$ , then  $\gamma(T) = (\omega, \sigma - A)$ , where  $\sigma = \text{diag}(\sigma_a)_{a \in S}$ .

### References.

- [BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, <http://www.math.toronto.edu/~drorbn/papers/KBH/>, arXiv:1308.1721.
- [BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects: From Alexander to Kashiwara and Vergne*, <http://www.math.toronto.edu/~drorbn/papers/WKO/>.
- [BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.
- [CR] A. S. Cattaneo and C. A. Rossi, *Wilson Surfaces and Higher Dimensional Knot Invariants*, Commun. in Math. Phys. **256-3** (2005) 513–537, arXiv:math-ph/0210037.
- [CT] D. Cimasoni and V. Turaev, *A Lagrangian Representation of Tangles*, Topology **44** (2005) 747–767, arXiv:math.GT/0406269.
- [KLW] P. Kirk, C. Livingston, and Z. Wang, *The Gassner Representation for String Links*, Comm. Cont. Math. **3** (2001) 87–136, arXiv:math/9806035.
- [LD] J. Y. Le Dimet, *Enlacements d'Intervalles et Représentation de Gassner*, Comment. Math. Helv. **67** (1992) 306–315.


**Theorem 3 [BND, BN].**  $\exists!$  a homomorphic expansion, aka a homomorphic universal finite type invariant  $Z$  of w-knotted balloons and hoops.  $\zeta := \log Z$  takes values in  $FL(T)^H \times CW(T)$ .

$\zeta$  is computable!  $\zeta$  of the Borromean tangle, to degree 5:



I have a nice free-Lie calculator!  
 $\zeta = \sum c_i \zeta_i$

**Proposition [BN].** Modulo all relations that universally hold for the 2D non-Abelian Lie algebra and after some changes-of-variable,  $\zeta$  reduces to  $\beta$  and the KBH operations on  $\zeta$  reduce to the formulas in Theorem 2.



**A Big Question.** Does it all extend to arbitrary 2-knots (not necessarily “simple”)? To arbitrary codimension-2 knots?

**BF Following [CR].**  $A \in \Omega^1(M = \mathbb{R}^4, \mathfrak{g})$ ,  $B \in \Omega^2(M, \mathfrak{g}^*)$ ,

$$S(A, B) := \int_M \langle B, F_A \rangle.$$

With  $\kappa: (S = \mathbb{R}^2) \rightarrow M$ ,  $\beta \in \Omega^0(S, \mathfrak{g})$ ,  $\alpha \in \Omega^1(S, \mathfrak{g}^*)$ , set

$$O(A, B, \kappa) := \int \mathcal{D}\beta \mathcal{D}\alpha \exp \left( \frac{i}{\hbar} \int_S \langle \beta, d_\kappa A \alpha + \kappa^* B \rangle \right).$$




Rossi

**The BF Feynman Rules.** For an edge  $e$ , let  $\Phi_e$  be its direction, in  $S^3$  or  $S^1$ . Let  $\omega_3$  and  $\omega_1$  be volume forms on  $S^3$  and  $S^1$ . Then

$$Z_{BF} = \sum_{\text{diagrams } D} \frac{[D]}{|\text{Aut}(D)|} \int_{S^2} \dots \int_{S^2} \int_{\mathbb{R}^4} \dots \int_{\mathbb{R}^4} \prod_{e \in D} \Phi_e^* \omega_3 \prod_{e \in D} \Phi_e^* \omega_1$$

(modulo some  $STU$ - and  $IHX$ -like relations).



**Issues.** • Signs don't quite work out, and BF seems to reproduce only “half” of the wheels invariant.

• There are many more configuration space integrals than BF Feynman diagrams and than just trees and wheels.

• I don't know how to define “finite type” for arbitrary 2-knots.



“God created the knots, all else in topology is the work of mortals.”

Leopold Kronecker (modified)



[www.katlas.org](http://www.katlas.org)

Add a quick combinatorial }?