

M.Sc. Math Workshop — Assignment #2

HUJI Spring 1998

Dror Bar-Natan

- (7) Prove: If $\lambda > 0$ is irrational then there exists a continuous function $\phi : [0, 1] \rightarrow [0, 1]$ so that for every $\epsilon > 0$ and every every continuous function $f : [0, 1] \times [0, 1] \rightarrow \mathbf{R}$ there exists a continuous function $g : [0, 1 + \lambda] \rightarrow \mathbf{R}$ so that $|f(x, y) - g(\phi(x) + \lambda\phi(y))| < \epsilon$ on a set of area at least $1 - \epsilon$ in $[0, 1] \times [0, 1]$. (Notice the different order of the quantifiers relative to Q2).
- (8) Can you completely cover a disk of diameter 100 with 99 rectangles of sides 100×1 ?
- (9) Let A and B be two $n \times n$ matrices over \mathbf{C} .
- (a) Prove that $[A, B] \stackrel{\text{def}}{=} AB - BA \neq I$.
 - (b) Prove that if $[A, [A, B]] = 0$, then $[A, B]$ is nilpotent.
- (10) Find a 2-variable polynomial that is always positive on \mathbf{R}^2 , and has exactly two critical points, both of which are minima.
- (11) Let $C_n = \{(z_1, \dots, z_n) \in \mathbf{C}^n : \forall 1 \leq j \leq n, z_i \neq z_j\}$ be the configuration space of n distinct points in the complex plane \mathbf{C} , and let the n th pure braid group $PB_n = \pi_1(C_n)$ be the fundamental group of C_n . Prove that $PB_n \simeq PB_{n-1} \rtimes F_{n-1}$ where F_{n-1} denotes the free group on $n - 1$ generators. Deduce that $PB_n \simeq F_1 \rtimes F_2 \rtimes \dots \rtimes F_{n-1}$.
- (12) Let $A_n = \{1, 2, 3\}^n$ and let $A = A_1$. A function $f : A_n \rightarrow A$ is called *injective*, if whenever x and y are different, $f(x) \neq f(y)$. Obviously, injective functions exist only if $n = 1$, and in this case, they are simply permutations $\pi : A \rightarrow A$. A function $f : A_n \rightarrow A$ is called *weakly injective*, if whenever x and y are *totally different*, meaning that $x_i \neq y_i$ for *all* $1 \leq i \leq n$, one has $f(x) \neq f(y)$. Prove that if $f : A_n \rightarrow A$ is weakly injective then for some permutation $\pi : A \rightarrow A$ and $1 \leq i \leq n$, one has $f(x) = \pi(x_i)$ for all $x \in A_n$.
- (13) f is a real valued function on the reals, and it is known that at any point at least one derivative of f vanishes (possibly different derivatives at different points). Prove that f is a polynomial.
- (14) What is the configuration space of the machine M_3 in the picture below?




Figure 3. The machine M_3 .