

The Variables:

$$\begin{pmatrix} b_1 \setminus a_1 & x_{12} & x_{13} \\ y_{12} & b_2 \setminus a_2 & x_{23} \\ y_{13} & y_{23} & b_3 \setminus a_3 \end{pmatrix}, \quad \xi_-, \alpha_-, \beta_-, \eta_-.$$

We short “ x_1 ” for either of x_{12}, x_{23} and “ x_2 ” for x_{13} . Weights are intuitive on yb and 3-complementary on ax : wt: $b \rightarrow 0, y_1 \rightarrow 1, y_2 \rightarrow 2, a \rightarrow 3, x_1 \rightarrow 2, x_2 \rightarrow 1$. Weights are 3-complementary on the dual (greek) variables.

In $m[ij \rightarrow k]$:

At $\epsilon = 0$: $\dots, \xi_1 \eta_1 b, \xi_2 \eta_2 b, \xi_1' \xi_1'' x_2, \eta_2 \xi_1 y_1, \alpha \xi_i x_i \dots$

At ϵ/ϵ^2 : $\dots, \xi_1 \eta_1 a, \xi_2 \eta_2 a, \dots$

In $\Delta[i \rightarrow jk]$:

At $\epsilon = 0$: $\dots, \eta_i b y_i, \dots$