




**Abstract.** There is expected to be a hidden paradise of poly-time computable knot polynomials lying just beyond the Alexander polynomial. I will describe my brute attempts to gain entry.

For long knots,  $\omega$  is Alexander, and that's the fastest Alexander algorithm I know! Dunfield: 1000-crossing fast.



**Why "expected"?** Gauss diagram  $v_{d,f}(K) = \sum_{Y \subset X(K), |Y|=d} f(Y)$  formulas [PV, GPV] show that finite-type invariants are all poly-time, and tempt to conjecture that there are no others. But Alexander shows it nonsense:

**Theorem** [EK, Ha, En, Se]. There is a "homomorphic expansion"



$d$	2	3	4	5	6	7	8	...
known invts* in $O(n^d)$	1	1	$\infty$	3	4	8	11	...

This is an unreasonable picture! \*Fresh, numerical, no cheating. So there ought to be further poly-time invariants.

(it is enough to know  $Z$  on  $\bowtie$  and have disjoint union and stitching formulas) ... exponential and too hard!

**Also.** • The diagonal above the Alexander diagonal in the Melvin-Morton-Rozansky [MM, Ro] of the coloured Jones polynomial. • The 2-loop contribution to the Kontsevich integral.

**Idea.** Look for "ideal" quotients of  $\mathcal{A}_n^v$  that have poly-sized descriptions; ... specifically, limit the co-brackets.

**Why "paradise"?** Foremost answer: **OBVIOUSLY.** Cf. proving (incomputable A)=(incomputable B), or categorifying (incomputable C).

**1-co and 2-co, aka TC and  $TC^2$** , on right. The primitives that remain are:

ωβ/K17: (extend to tangles, perhaps detect non-slice ribbon knots)




Figure 1. A tangle.




Figure 2. A ribbon singularity, a clasp singularity, and an example of a ribbon knot.

**Second Answer.** The second answer has to do with "Algebraic Knot Theory", so let me start with that. Somewhat informally, a "tangle" is a piece of a knot, or a "knot with endpoints" (an example is on the right). Knots can be assembled by stitching together the strands of several tangles, or the different strands of a single tangle. Some interesting classes of knots can be defined algebraically using tangles and these stitching operations. Here is the most interesting example:  
**Definition 1.** A "ribbon knot" is a knot  $K$  that can be presented as the boundary of a disk  $D$  which is allowed to have "ribbon singularities" but not "clasp singularities". See Figure 2.  
**Definition 2.** Let  $\mathcal{T}_n$  denote the set of all tangles  $T$  with  $2n$  components that connect  $2n$  points along a "top end" with  $2n$  points along a "bottom end" inducing the identity permutation of ends (an example is the tangle in Figure 1). Given  $T \in \mathcal{T}_n$ , let  $\tau(T)$  be the result of stitching its components at the top in pairs as on the right — it is an  $n$ -component tangle all of whose ends are at the bottom, and we (somewhat loosely) denote the set of all such by  $\mathcal{T}_n$ , so  $\tau: \mathcal{T}_n \rightarrow \mathcal{T}_n$ . Likewise let  $\sigma(T)$  be the result of stitching  $T$  both at the top and at the bottom. Also as on the right. So  $\sigma(T)$  is a 1-component tangle, which is the same as a knot, and  $\sigma: \mathcal{T}_n \rightarrow \mathcal{T}_1$ .  
**Theorem 1** (I have not seen this theorem in the literature, yet it is not difficult to prove). The set of ribbon knots is the set of all knots  $K$  that can be written as  $K = \sigma(T)$  for some tangle  $T$  for which  $\tau(T)$  is the untagged (crossingless) tangle  $U$ :  
 (ribbon knots) =  $\{\sigma(T) : T \in \mathcal{T}_n \text{ and } \tau(T) = U \in \mathcal{T}_n\}$ .

Now suppose we have an invariant  $Z: \mathcal{T}_1 \rightarrow A_1$  of tangles, which takes values in some spaces  $A_i$ . Suppose also we have operations  $\tau_1: A_2 \rightarrow A_1$  and  $\sigma_1: A_2 \rightarrow A_1$  such that the diagram on the right is commutative. Then

$$Z(\text{ribbon knots}) \subseteq \mathcal{R}_1 := \{k_1(C) : C \in A_2 \text{ and } \tau_1(C) = \sigma_1(C) \in A_1\} \quad (2)$$

where  $\tau_1 := Z(U) \in A_1$ . If the target spaces  $A_i$  are algebraic (polynomials, matrices, matrices of polynomials, etc.) and the operations  $\tau_1$  and  $\sigma_1$  are algebraic maps between them (at this stage, meaning just "have simple algebraic formulas"), then  $\mathcal{R}_1$  is an algebraically defined set. Hence we potentially have an algebraic way to detect non-ribbon knots: if  $Z(K) \notin \mathcal{R}_1$ , then  $K$  is not ribbon.

As it turns out, it is valuable to detect non-ribbon knots. Indeed the Slice-Ribbon Conjecture (Fox, 1960s) asserts that every slice knot (a knot in  $S^3$  that can be presented as the boundary of a disk embedded in  $B^4$ ) is ribbon. Gompf, Scharlemann, and Thompson [GST] describe a family of slice knots which they conjecture are not ribbon (the simplest of those is on the right). With the algebraic technology described above it may be possible to show that the [GST] knots are indeed non-ribbon, thus disproving the Slice-Ribbon Conjecture.

- What would it take?
- C1. An invariant  $Z$  which makes sense on tangles and for which diagram (1) commutes.
  - C2.  $Z$  cannot be a simple extension of the Alexander polynomial to tangles, for by Fox-Milnor [FM] the Alexander polynomial does not detect non-ribbon slice knots.
  - C3.  $Z$  cannot be computable from finitely many finite type invariants, for this would contradict the results of Ng [Ng].
  - C4.  $Z$  must be computable on at least the simplest [GST] knot, which has 48 crossings.
  - C5. It is better if in some meaningful sense the size of the spaces  $A_i$  grows slowly in  $k$ . Indeed, if  $A_{2k}$  is much bigger than  $A_k$  and  $A_1$  then at least generally  $\mathcal{R}_k$  will be the full set  $A_k$  and our condition will be empty.

No invariant that I know now meets these criteria. Alexander and Vassiliev fail C2 and C3, respectively. Almost all quantum invariants and knot homologies pass C1-C3, but fail C4. Jones, HOMFLY-PT and Khovanov potentially pass C4, yet fail C5. We must come up with something new.

[FM] R. H. Fox and J. W. Milnor, *Singularities of 2-Spheres in 4-Space and Cobordism of Knots*, Osaka J. Math. **3** (1966) 257–267.  
 [GST] R. E. Gompf, M. Scharlemann, and A. Thompson, *Fibered Knots and Potential Counterexamples to the Property 2R and Slice-Ribbon Conjecture*, Geom. and Top. **14** (2010) 2305–2347, arXiv:1103.1601.  
 [Ng] K. Y. Ng, *Groups of ribbon knots*, Topology **37** (1998) 441–458, arXiv:q-alg/9502017 (with an addendum at arXiv:math.GT/0310074)

<sup>1</sup>A slight subtlety arises: There is no taking limits here, and C3 does not preclude the possibility that  $Z$  is computable from infinitely many finite type invariants. The Fox-Milnor condition on the Alexander polynomial of ribbon knots, for example, is expressible in terms of the full Alexander polynomial, yet not in terms of any finite type reduction thereof. Unfortunately by C2 it cannot be used here.

**Why "brute"?** Cause it's the only thing I know, for now. There may be better ways in, and it's fair to hope that sooner or later they will be found.



**The Gold Standard** is set by the formulas [BNS, BN] for Alexander. An  $S$ -component tangle  $T$  has  $\Gamma(T) \in R_S \times M_{S \times S}(R_S) = \left\{ \begin{matrix} \omega & S \\ S & A \end{matrix} \right\}$  with  $R_S := \mathbb{Z}\langle\{t_a : a \in S\}\rangle$ :

$$\left( \begin{matrix} a & b \\ a & b \end{matrix} \right) \rightarrow \begin{matrix} a & b \\ a & 1 - t_a^{\pm 1} \\ b & 0 \\ b & t_a^{\pm 1} \end{matrix} \quad T_1 \sqcup T_2 \rightarrow \begin{matrix} \omega_1 \omega_2 & S_1 & S_2 \\ S_1 & A_1 & 0 \\ S_2 & 0 & A_2 \end{matrix}$$

$$\begin{matrix} \omega & a & b & S \\ a & \alpha & \beta & \theta \\ b & \gamma & \delta & \epsilon \\ S & \phi & \psi & \Xi \end{matrix} \rightarrow \begin{matrix} m_c^{ab} \\ t_a, t_b \rightarrow t_c \\ \mu := 1 - \beta \end{matrix} \quad \left( \begin{matrix} \mu \omega & c & S \\ c & \gamma + \alpha \delta / \mu & \epsilon + \delta \theta / \mu \\ S & \phi + \alpha \psi / \mu & \Xi + \psi \theta / \mu \end{matrix} \right)$$

**The 2D relations** come from the relation with 2D Lie bialgebras:



TODO: a schematic picture of stitching

TODO: The 2D relations.

**References.**

[BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, ωβ/KBH, arXiv:1308.1721.  
 [BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I, II, IV*, ωβ/WK01, ωβ/WK02, ωβ/WK04, arXiv:1405.1956, arXiv:1405.1955, arXiv:1511.05624.  
 [BNG] D. Bar-Natan and S. Garoufalidis, *On the Melvin-Morton-Rozansky conjecture*, Invent. Math. **125** (1996) 103–133.  
 [BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.  
 [En] B. Enriquez, *A Cohomological Construction of Quantization Functors of Lie Bialgebras*, Adv. in Math. **197-2** (2005) 430–479, arXiv:math/0212325.  
 [EK] P. Etingof and D. Kazhdan, *Quantization of Lie Bialgebras, I*, Selecta Mathematica **2** (1996) 1–41, arXiv:q-alg/9506005.  
 [GPV] M. Goussarov, M. Polyak, and O. Viro, *Finite type invariants of classical and virtual knots*, Topology **39** (2000) 1045–1068, arXiv:math.GT/9810073.  
 [Ha] A. Haviv, *Towards a diagrammatic analogue of the Reshetikhin-Turaev link invariants*, Hebrew University PhD thesis, Sep. 2002, arXiv:math.QA/0211031.  
 [MM] P. M. Melvin and H. R. Morton, *The coloured Jones function*, Commun. Math. Phys. **169** (1995) 501–520.  
 [PV] M. Polyak and O. Viro, *Gauss Diagram Formulas for Vassiliev Invariants*, Inter. Math. Res. Notices **11** (1994) 445–453.  
 [Ro] L. Rozansky, *A contribution of the trivial flat connection to the Jones polynomial and Witten's invariant of 3d manifolds, I*, Comm. Math. Phys. **175-2** (1996) 275–296, arXiv:hep-th/9401061.  
 [Se] P. Ševera, *Quantization of Lie Bialgebras Revisited*, Sel. Math., NS, to appear, arXiv:1401.6164.



“God created the knots; all else in topology is the work of mortals.”

Leopold Kronecker (modified)



[katlas.org](http://katlas.org)