

[BN2] D. Bar-Natan, *Finite Type Invariants*, in *Encyclopedia of Mathematical Physics*, (J.-P. Francoise, G. L. Naber and Tsou S. T., eds.) Elsevier, Oxford, 2006 (vol. 2 p. 340).

[Book] The reviewed book.

[BL] J. S. Birman and X.-S. Lin, *Knot polynomials and Vassiliev's invariants*, *Invent. Math.* **111** (1993) 225–270.

[Cv] P. Cvitanović, *Group Theory, Birdtracks, Lie's, and Exceptional Groups*, Princeton University Press, Princeton 2008 and <http://www.birdtracks.eu>.

[Go1] M. Goussarov, *A new form of the Conway-Jones polynomial of oriented links*, *Zapiski nauch. sem. POMI* **193** (1991) 4–9 (English translation in *Topology of manifolds and varieties* (O. Viro, editor), Amer. Math. Soc., Providence 1994, 167–172).

[Go2] M. Goussarov, *On n-equivalence of knots and invariants of finite degree*, *Zapiski nauch. sem. POMI* **208** (1993) 152–173 (English translation in *Topology of manifolds and varieties* (O. Viro, editor), Amer. Math. Soc., Providence 1994, 173–192).

[Jo] V. F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, *Bull. Amer. Math. Soc.* **12** (1985) 103–111.

[Ko1] M. Kontsevich, *Vassiliev's knot invariants*, *Adv. in Sov. Math.*, **16(2)** (1993) 137–150.

[Ko2] M. Kontsevich, *Feynman diagrams and low-dimensional topology*, First European Congress of Mathematics **II** 97–121, Birkhäuser Basel 1994.

[Pe] R. Penrose, *Applications of negative dimensional tensors*, *Combinatorial mathematics and its applications* (D. J. A. Welsh, ed.), Academic Press, San-Diego 1971, 221–244.

[RT] N. Yu. Reshetikhin and V. G. Turaev, *Ribbon graphs and their invariants derived from quantum groups*, *Commun. Math. Phys.* **127** (1990) 1–26.

[Va1] V. A. Vassiliev, *Cohomology of knot spaces*, in *Theory of Singularities and its Applications (Providence)* (V. I. Arnold, ed.), Amer. Math. Soc., Providence, 1990.

[Va2] V. A. Vassiliev, *Complements of discriminants of smooth maps: topology and applications*, *Trans. of Math. Mono.* **98**, Amer. Math. Soc., Providence, 1992.

[Wi] E. Witten, *Quantum field theory and the Jones polynomial*, *Commun. Math. Phys.* **121** (1989) 351–399.

Dror Bar-Natan

University of Toronto, Canada

December 6, 2019 (first edition February 7, 2013)

My talk yesterday:

More Dror: $\omega\epsilon\beta$ /talks

Dror Bar-Natan: Talks: Toronto-1912 [oeβ:=http://drorbn.net/to19/](http://drorbn.net/to19/)

Geography vs. Identity
Thanks for inviting me to the *Topology* session!

Abstract. Which is better, an emphasis on where things happen or on who are the participants? I can't tell; there are advantages and disadvantages either way. Yet much of quantum topology seems to be heavily and unfairly biased in favour of geography.

Geographers care for placement; for them, braids and tangles have ends at some distinguished points, hence they form categories whose objects are the placements of these points. For them, the basic operation is a binary "stacking of tangles". They are lead to monoidal categories, braided monoidal categories, representation theory, and much or most of we call "quantum topology".

Identifiers believe that strand identity persists even if one crosses or is being crossed. The key operation is a unary stitching operation m_c^{ab} , and one is lead to study meta-monoids, meta-Hopf-algebras, etc. See oeβ/leg, oeβ/kbh.

Braids.

Geography: $GB := \langle \gamma_i \mid \gamma_i \gamma_k = \gamma_k \gamma_i \text{ when } |i-k| > 1 \rangle = B$.
(captures quantum algebra!)

Identity: $IB := \langle \sigma_{ij} \mid \sigma_{ij} \sigma_{kl} = \sigma_{kl} \sigma_{ij} \text{ when } \{|i, j, k, l\} = 4\} = PAB$.

Theorem. Let $S = \langle \tau \rangle$ be the symmetric group. Then \mathfrak{B} is both $PAB \rtimes S \cong B * S \langle \gamma_i \tau = \tau \gamma_i \text{ when } \tau i = j, \tau(i+1) = (j+1) \rangle$ (and so PAB is "bigger" than B , and hence quantum algebra doesn't see topology very well).

Proof. Going left, $\gamma_i \mapsto \sigma_{i,i+1}(i+1)$. Going right, if $i < j$ map $\sigma_{ij} \mapsto (j-1 \ j-2 \ \dots \ i) \gamma_{j-1}(i+1 \ \dots \ j)$ and if $i > j$ use $\sigma_{ij} \mapsto (j \ j+1 \ \dots \ i) \gamma_j(i-1 \ \dots \ j+1)$.

\mathfrak{B} views of σ_{ij} :

The Burau Representation of PAB_n acts on $\mathbb{R}^n := \mathbb{Z}[t^{\pm 1}]^n = R(v_1, \dots, v_n)$ by $\sigma_{ij} v_k = v_k + \delta_{kj}(t-1)(v_j - v_i)$.

$\delta := \delta_{i,j} := \mathbf{1f}[i=j, 1, 0]$ oeβ/code
 $\mathfrak{B}_{i,j}[\mathcal{L}] := \mathcal{L} / \cdot v_k \mapsto v_k + \delta_{kj}(t-1)(v_j - v_i) // \text{Expand}$
 $(\text{bas3} = \{v_1, v_2, v_3\} // \mathfrak{B}_{1,2}) // \mathfrak{B}_{1,2}$
 $\{v_{1,1} v_1 - t v_{1,1} + t v_{2,1}\} // \mathfrak{B}_{1,2}$
 $\text{bas3} // \mathfrak{B}_{1,2} // \mathfrak{B}_{1,3} // \mathfrak{B}_{2,3}$
 $\{v_{1,1} v_1 - t v_{1,1} + t v_{2,1}, v_{1,1} - t v_{1,1} + t v_{2,1} - t^2 v_{2,1} + t^2 v_{3,1}\}$
 $\text{bas3} // \mathfrak{B}_{2,3} // \mathfrak{B}_{1,3} // \mathfrak{B}_{1,2}$
 $\{v_{1,1} v_1 - t v_{1,1} + t v_{2,1}, v_{1,1} - t v_{1,1} + t v_{2,1} - t^2 v_{2,1} + t^2 v_{3,1}\}$

S_n acts on \mathbb{R}^n by permuting the v_i , so the Burau representation extends to \mathfrak{B}_n and restricts to B_n . With this, γ_i maps $v_i \mapsto v_{i+1}, v_{i+1} \mapsto v_i(1-t)v_{i+1}$, and otherwise $v_k \mapsto v_k$.

Geography view:
 $\gamma_1 = \times \mid \mid \mid \quad \gamma_2 = \mid \times \mid \mid \quad \gamma_3 = \mid \mid \times \dots$
so x is γ_2 .

Identity view:
At x strand 1 crosses strand 3, so x is σ_{13} .

The Gold Standard is set by the "T-calculus" Alexander formulas (oeβ/mac). An S -component tangle T has $\Gamma(T) \in R_S \times M_{S \times S}(R_S) = \begin{pmatrix} \omega & S \\ S & A \end{pmatrix}$ with $R_S := \mathbb{Z}\langle T_a : a \in S \rangle$:

$(a' \times, b' \times, a) \rightarrow \frac{1}{a} \mid \frac{1}{b} \mid \frac{1-t}{T_a} \mid \frac{\omega_1 \omega_2}{S_1} \mid \frac{S_1}{S_2} \mid \frac{S_1}{A_1} \mid \frac{0}{A_2}$

$\omega \mid \frac{a}{\alpha} \mid \frac{b}{\beta} \mid \frac{S}{\theta} \mid \frac{m_c^{ab}}{\delta} \mid \frac{c}{S} \mid \frac{(1-\beta)\omega}{\phi + \frac{\alpha\beta}{1-\beta}} \mid \frac{c}{S} \mid \frac{S}{\Xi + \frac{\alpha\beta}{1-\beta}}$

$\frac{a}{b} \mid \frac{\alpha}{\gamma} \mid \frac{\beta}{\delta} \mid \frac{S}{\phi} \mid \frac{m_c^{ab}}{\Xi} \mid \frac{c}{S} \mid \frac{(1-\beta)\omega}{\phi + \frac{\alpha\beta}{1-\beta}} \mid \frac{c}{S} \mid \frac{S}{\Xi + \frac{\alpha\beta}{1-\beta}}$

$T_a \mid T_b \rightarrow T_c$

The Gassner Representation of PAB_n acts on $V = \mathbb{R}^n := \mathbb{Z}[t^{\pm 1}]^n = R(v_1, \dots, v_n)$ by $\sigma_{ij} v_k = v_k + \delta_{kj}(t-1)(v_j - v_i)$.

$\mathfrak{G}_{i,j}[\mathcal{L}] := \mathcal{L} / \cdot v_k \mapsto v_k + \delta_{kj}(t-1)(v_j - v_i) // \text{Expand}$
 $(\text{bas3} // \mathfrak{G}_{1,2} // \mathfrak{G}_{1,3} // \mathfrak{G}_{2,3}) = (\text{bas3} // \mathfrak{G}_{1,2} // \mathfrak{G}_{1,3} // \mathfrak{G}_{2,3})$

True
 S_n acts on \mathbb{R}^n by permuting the v_i and the t_i , so the Gassner representation extends to \mathfrak{B}_n and then restricts to B_n as a \mathbb{Z} -linear ∞ -dimensional representation. It then descends to PB_n as a finite-rank R -linear representation, with lengthy non-local formulas.

Geographers: Gassner is an obscure partial extension of Burau.
Identifiers: Burau is a trivial silly reduction of Gassner.

The Turbo-Gassner Representation. With the same R and V , TG acts on $V \oplus (R^n \oplus \mathfrak{G}) \oplus (S^2 V \oplus V^n) = R(v_i, u_i, u_i u_j, w_i)$ by $\mathfrak{TG}_{i,j}[\mathcal{L}] := \mathcal{L} / \cdot \{$

$v_k \mapsto v_k + \delta_{kj}((t-1)(v_j - v_i) + v_{i,j} - v_{i,i}) + \delta_{ki}(u_j - u_i) u_i w_j,$
 $v_{i,i} \mapsto v_{i,i} + (t-1) \cdot$
 $(\delta_{i,j}(v_{i,j} - v_{i,i}) + (\delta_{i,i} - \delta_{i,j} t^2) t_j)$
 $(u_i + \delta_{i,j}(t-1)(u_j - u_i)) u_i w_j,$
 $u_i \mapsto u_i + \delta_{i,j}(t-1)(u_j - u_i),$
 $w_i \mapsto w_i + (\delta_{i,j} - \delta_{i,i})(t^2 - 1) w_i // \text{Expand}$
 $\text{bas3} = \{v_1, v_2, v_3, u_1^2 w_1, u_1^2 w_2, u_1^2 w_3, v_{1,1}, v_{2,1}, v_{3,1}, v_{1,2}, v_{2,2}, v_{3,2}, v_{1,3}, v_{2,3}, u_1^2 w_1, u_1^2 w_2, u_1^2 w_3, u_1 u_2 w_1, u_1 u_2 w_2, u_1 u_2 w_3, u_2 u_1 w_1, u_2 u_1 w_2, u_2 u_1 w_3, u_2 u_2 w_1, u_2 u_2 w_2, u_2 u_2 w_3, u_2 u_3 w_1, u_2 u_3 w_2, u_2 u_3 w_3\};$
 $(\text{bas3} // \mathfrak{TG}_{1,2} // \mathfrak{TG}_{1,3} // \mathfrak{TG}_{2,3}) = (\text{bas3} // \mathfrak{TG}_{1,2} // \mathfrak{TG}_{1,3} // \mathfrak{TG}_{2,3})$

True
Like Gassner, TG is also a representation of PB_n .

I have no idea where it belongs!

My talk tomorrow, at the *chord diagrams everywhere* session: oeβ/talks

Picture credits: Rope from "The Project Gutenberg eBook, Knots, Splices and Rope Work, by A. Hyatt Verrill", <http://www.gutenberg.org/files/13510/13510-h/13510-h.htm>. Plane from NASA, <http://www.grc.nasa.gov/WWW/k-12/airplane/rotations.html>.