


**Theorem 2 [BND].**  $\exists!$  a homomorphic expansion, aka a homomorphic universal finite type invariant  $Z^w$  of pure w-tangles.  $z^w := \log Z^w$  takes values in  $FL(S)^S \times CW(S)$ .

$z$  is computable.  $z$  of the Borromean tangle, to degree 5 [BN]:



**Definition.** (Compare [BNS, BN]) A **The Abstract Context** meta-monoid is a functor  $M: (\text{finite sets, injections}) \rightarrow (\text{sets})$  (think “ $M(S)$  is quantum  $G^S$ ”, for  $G$  a group) along with natural operations  $*$ :  $M(S_1) \times M(S_2) \rightarrow M(S_1 \sqcup S_2)$  whenever  $S_1 \cap S_2 = \emptyset$  and  $m_c^{ab}: M(S) \rightarrow M((S \setminus \{a, b\}) \sqcup \{c\})$  whenever  $a \neq b \in S$  and  $c \notin S \setminus \{a, b\}$ , such that

$$\text{meta-associativity: } m_a^{ab} // m_a^{ac} = m_b^{bc} // m_a^{ab}$$

$$\text{meta-locality: } m_c^{ab} // m_f^{de} = m_f^{de} // m_c^{ab}$$

and, with  $\epsilon_b = M(S \hookrightarrow S \sqcup \{b\})$ ,

$$\text{meta-unit: } \epsilon_b // m_a^{ab} = Id = \epsilon_b // m_a^{ba}$$

**Claim.** Pure virtual tangles  $PT$  form a meta-monoid.

**Theorem.**  $S \mapsto \Gamma_0(S)$  is a meta-monoid and  $z_0: PT \rightarrow \Gamma_0$  is a morphism of meta-monoids.

**Strong Conviction.** There exists an extension of  $\Gamma_0$  to a bigger meta-monoid  $\Gamma_{01}(S) = \Gamma_0(S) \times \Gamma_1(S)$ , along with an extension of  $z_0$  to  $z_{01}: PT \rightarrow \Gamma_{01}$ , with

$$\Gamma_1(S) = V \oplus V^{\otimes 2} \oplus V^{\otimes 3} \oplus S^2(V)^{\otimes 2} \quad (\text{with } V := R_S(S)).$$


Furthermore, upon reducing to a single variable everything is polynomial size and polynomial time.

Furthermore,  $\Gamma_{01}$  is given using a “meta-2-cocycle  $\rho_c^{ab}$  over  $\Gamma_0$ ”: In addition to  $m_c^{ab} \rightarrow m_{0c}^{ab}$ , there are  $R_S$ -linear  $m_{1c}^{ab}: \Gamma_1(S \sqcup \{a, b\}) \rightarrow \Gamma_1(S \sqcup \{c\})$ , a meta-right-action  $\alpha^{ab}: \Gamma_1(S) \times \Gamma_0(S) \rightarrow \Gamma_1(S)$   $R_S$ -linear in the first variable, and a first order differential operator (over  $R_S$ )  $\rho_c^{ab}: \Gamma_0(S \sqcup \{a, b\}) \rightarrow \Gamma_1(S \sqcup \{c\})$  such that



$$(\zeta_0, \zeta_1) // m_c^{ab} = (\zeta_0 // m_{0c}^{ab}, (\zeta_1, \zeta_0) // \alpha^{ab} // m_{1c}^{ab} + \zeta_0 // \rho_c^{ab})$$

**What's done?** The braid part, with still-ugly formulas.

**What's missing?** A lot of concept- and detail-sensitive work towards  $m_{1c}^{ab}$ ,  $\alpha^{ab}$ , and  $\rho_c^{ab}$ . The “ribbon element”.




Back to v – the 2D “Jones Quotient”.



**References.**

[ADO] Y. Akutsu, T. Deguchi, and T. Ohtsuki, *Invariants of Colored Links*, J. of Knot Theory and its Ramifications **1-2** (1992) 161–184.  
 [BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*,  $\omega\epsilon\beta$ /KBH, arXiv:1308.1721.  
 [BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I-II*,  $\omega\epsilon\beta$ /WKO1,  $\omega\epsilon\beta$ /WKO2, arXiv:1405.1956, arXiv:1405.1955.  
 [BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.  
 [CT] D. Cimasoni and V. Turaev, *A Lagrangian Representation of Tangles*, Topology **44** (2005) 747–767, arXiv:math.GT/0406269.  
 [En] B. Enriquez, *A Cohomological Construction of Quantization Functors of Lie Bialgebras*, Adv. in Math. **197-2** (2005) 430–479, arXiv:math/0212325.  
 [EK] P. Etingof and D. Kazhdan, *Quantization of Lie Bialgebras, I*, Selecta Mathematica **2** (1996) 1–41, arXiv:q-alg/9506005.  
 [GST] R. E. Gompf, M. Scharlemann, and A. Thompson, *Fibered Knots and Potential Counterexamples to the Property 2R and Slice-Ribbon Conjectures*, Geom. and Top. **14** (2010) 2305–2347, arXiv:1103.1601.  
 [KLW] P. Kirk, C. Livingston, and Z. Wang, *The Gassner Representation for String Links*, Comm. Cont. Math. **3** (2001) 87–136, arXiv:math/9806035.  
 [LD] J. Y. Le Dimet, *Enlacements d'Intervalles et Représentation de Gassner*, Comment. Math. Helv. **67** (1992) 306–315.



**A bit about ribbon knots.** A “ribbon knot” is a knot that can be presented as the boundary of a disk that has “ribbon singularities”, but no “clasp singularities”. A “slice knot” is a knot in  $S^3 = \partial B^4$  which is the boundary of a non-singular disk in  $B^4$ . Every ribbon knot is clearly slice, yet,

**Conjecture.** Some slice knots are not ribbon.

**Fox-Milnor.** The Alexander polynomial of a ribbon knot is always of the form  $A(t) = f(t)f(1/t)$ . (also for slice)

[GST]: a slice knot that might not be ribbon (48 crossings).

