






Finite Type Invariants of Ribbon Knotted Balloons and Hoops

Abstract. On my September 17 Geneva talk (ω/sep) I described a certain trees-and-wheels-valued invariant ζ of ribbon knotted loops and 2-spheres in 4-space, and my October 8 Geneva talk (ω/oct) describes its reduction to the Alexander polynomial. Today I will explain how that same invariant arises completely naturally within the theory of finite type invariants of ribbon knotted loops and 2-spheres in 4-space.



My goal is to tell you why such an invariant is expected, yet not to derive the computable formulas.



Expansions
the semi-virtual $\otimes := \begin{matrix} \diagup \\ \diagdown \end{matrix} - \begin{matrix} \diagdown \\ \diagup \end{matrix}$ i.e. $\begin{matrix} \diagup \\ \diagdown \end{matrix} - \begin{matrix} \diagdown \\ \diagup \end{matrix}$ or $\begin{matrix} \diagdown \\ \diagup \end{matrix} - \begin{matrix} \diagup \\ \diagdown \end{matrix}$

Let $\mathcal{I}^n := \langle \text{pictures with } \geq n \text{ semi-virts} \rangle \subset \mathcal{K}^{bh}$.
We seek an "expansion"

$$Z: \mathcal{K}^{bh} \rightarrow \text{gr } \mathcal{K}^{bh} = \bigoplus_{n \geq 0} \mathcal{I}^n / \mathcal{I}^{n+1} =: \mathcal{A}^{bh}$$

satisfying "property U": if $\gamma \in \mathcal{I}^n$, then


$$Z(\gamma) = (0, \dots, 0, \gamma / \mathcal{I}^{n+1}, *, *, \dots).$$



X.-S. Lin

Why? • Just because, and this is vastly more general.
• $(\mathcal{K}^{bh} / \mathcal{I}^{n+1})^*$ is "finite-type/polynomial invariants".
• The Taylor example: Take $\mathcal{K} = C^\infty(\mathbb{R}^n)$, $\mathcal{I} = \{f \in \mathcal{K} : f(0) = 0\}$. Then $\mathcal{I}^n = \{f : f \text{ vanishes like } |x|^n\}$ so $\mathcal{I}^n / \mathcal{I}^{n+1}$ is homogeneous polynomials of degree n and Z is a "Taylor expansion"! (So Taylor expansions are vastly more general than you'd think).


Plan. We'll construct a graded $\tilde{\mathcal{A}}^{bh}$, a surjective graded $\pi: \tilde{\mathcal{A}}^{bh} \rightarrow \mathcal{A}^{bh}$, and a filtered $\tilde{Z}: \mathcal{K}^{bh} \rightarrow \tilde{\mathcal{A}}^{bh}$ so that $\pi \parallel \text{gr } \tilde{Z} = Id$ (property U: if $\deg D = n$, $\tilde{Z}(\pi(D)) = \pi(D) + (\deg \geq n)$). Hence • π is an isomorphism. • $Z := \tilde{Z} \parallel \pi$ is an expansion.




"God created the knots, all else in topology is the work of mortals."

Leopold Kronecker (modified)


www.katlas.org




Action 1.




Deriving 4T.




Action 2.



The Bracket-Rise Theorem.



Proof.



Corollaries. (1) Related to Lie algebras! (2) Only trees and wheels persist.

Theorem. \mathcal{A}^{bh} is a bi-algebra. The space of its primitives is $FL(T)^H \times CW(T)$, and $\zeta = \log Z$.

ζ is computable! ζ of the Borromean tangle, to degree 5:

