

# Local Khovanov Homology (1)

(an outdated overview)


The Jones polynomial:

$$J : \text{link} \mapsto q \langle -q^2 \rangle, \quad J : \text{link} \mapsto -q^{-2} \langle -q^2 \rangle + q^{-1} \langle -q^2 \rangle$$

$$\bigcirc^k \mapsto (q + q^{-1})^k$$

$$J : \text{link} \mapsto -q^{-1} \langle -q^2 \rangle + \langle -q^2 \rangle + \langle -q^2 \rangle - q \langle -q^2 \rangle = -q^{-1} \langle -q^2 \rangle + (q + q^{-1}) \langle -q^2 \rangle - q \langle -q^2 \rangle$$

R2



Mikhail Khovanov


## What is it?

A cube for each knot/link projection;


Vertices: All fillings of




with




or with

Edges: All fillings of  $I \times$ with  $I \times$ 

or with




and precisely one




## Signs?

$$\begin{array}{c} dx \\ \swarrow \quad \searrow \\ dy \quad dz \\ \swarrow \quad \searrow \\ dx^2 dy^2 dz^2 \end{array}$$

## More crossings?



## General Crossings




## Where does it live?

In  $Kom(Mat(\langle Cob \rangle / \{S, T, G, NC\})) / \text{homotopy}$ 


Kom: Complexes Mat: Matrices

Cob: Cobordisms  $\langle \dots \rangle$ : Formal lin. comb.

Cob:



Mat(C):




$$S: \text{circle} = 0 \quad T: \text{torus} = 2 \quad G: \text{cylinder} = 0$$

$$NC: 2 \text{ (cylinder)} = \text{torus} + \text{cylinder} + \text{cylinder}$$

## Computable!

via

"complex simplification"



## Complexes:

$$\Omega = (\Omega^{-n} \longrightarrow \Omega^{-n+1} \longrightarrow \dots \longrightarrow \Omega^n)$$

## Morphisms:

$$\begin{array}{ccccccc} \dots & \longrightarrow & \Omega_0^{r-1} & \xrightarrow{d^{r-1}} & \Omega_0^r & \xrightarrow{d^r} & \Omega_0^{r+1} \longrightarrow \dots \\ & & \downarrow F^{r-1} & & \downarrow F^r & & \downarrow F^{r+1} \\ \dots & \longrightarrow & \Omega_1^{r-1} & \xrightarrow{d^{r-1}} & \Omega_1^r & \xrightarrow{d^r} & \Omega_1^{r+1} \longrightarrow \dots \end{array}$$

## Homotopies:

$$\begin{array}{ccccccc} \Omega_0^{r-1} & \xrightarrow{d^{r-1}} & \Omega_0^r & \xrightarrow{d^r} & \Omega_0^{r+1} \\ \downarrow F^{r-1} & \parallel G^{r-1} & \downarrow F^r & \parallel G^r & \downarrow F^{r+1} \\ \Omega_1^{r-1} & \xrightarrow{d^{r-1}} & \Omega_1^r & \xrightarrow{d^r} & \Omega_1^{r+1} \end{array}$$

$$F^r - G^r = h^{r+1} d^r + d^{r-1} h^r$$

**The Main Point.** "The cube",  $\text{Kh}(L)$ , is an up-to-homotopy invariant of knots and links. It's Euler characteristic is the Jones polynomial, yet it is strictly stronger than the Jones polynomial. It is functorial (in the appropriate sense) and practically computable.

**The Categorification Speculative Paradigm.** • Every object in math is the Euler characteristic of a complex.  
• Every operation lifts to an operation between complexes.  
• Every identity remains true, up to homotopy.

All arrows in an arbitrary additive category!