

A FAST, STRONG, TOPOLOGICALLY MEANINGFUL, AND FUN KNOT INVARIANT

DROR BAR-NATAN AND ROLAND VAN DER VEEN

ABSTRACT. In this paper we discuss a pair of polynomial knot invariants $\Theta = (\Delta, \theta)$ which is:

- Theoretically and practically fast: Θ can be computed in polynomial time. We can compute it in full on random knots with over 300 crossings, and its evaluation at simple rational numbers on random knots with over 600 crossings.
- Strong: Its separation power is much greater than the hyperbolic volume, the HOMFLY-PT polynomial and Khovanov homology (taken together) on knots with up to 15 crossings (while being computable on much larger knots).
- Topologically meaningful: It likely gives a genus bound, and there are reasons to hope that it would do more.
- Fun: Scroll to Figures 1.1–1.4, 3.1, and 6.2.

Δ is merely the Alexander polynomial. θ is almost certainly equal to an invariant that was studied extensively by Ohtsuki [Oh2], continuing Rozansky, Kricker, and Garoufalidis [Roz1, Roz2, Roz3, Kr, GR]. Yet our formulas, proofs, and programs are much simpler and enable its computation even on very large knots.

CONTENTS

1. Fun	2
2. The Main Theorem	4
3. Implementation and Examples	8
3.1. Implementation	8
3.2. Examples	10
4. Proof of the Main Theorem, Theorem 1	11
4.1. Proof of Invariance	11
4.2. Proof of Polynomiality	22
5. Strong and Meaningful	23
5.1. Strong	23
5.2. Meaningful	25
5.2.1. The Knot Genus	25
5.2.2. Fibered Knots	26
6. Stories, Conjectures, and Dreams	26
7. Acknowledgement	33
References	33

Date: First edition September 22, 2025. This edition December 11, 2025.

2020 Mathematics Subject Classification. Primary 57K14, secondary 16T99.

Key words and phrases. Alexander polynomial, loop expansion, solvable approximation, knot genus, fibered knots, ribbon knots, polynomial time computations, Feynman diagrams, perturbed Gaussian integration, Seifert formulas.


This paper is available in electronic form, along with source files and a demo *Mathematica* notebook at <http://drorbn.net/Theta> and at [arXiv:2509.18456](https://arxiv.org/abs/2509.18456).

1. FUN

The word “fun” rarely appears in the title of a math paper, so let us start with a brief justification.

Θ is a pair of polynomials. The first, Δ , is old news, the Alexander polynomial [Al]. It is a one-variable Laurent polynomial in a variable T . For example, $\Delta(\textcircled{3}) = T^{-1} - 1 + T$. We turn such a polynomial into a list of coefficients (for $\textcircled{3}$, it is $(1, -1, 1)$), and then to a chain of bars of varying colours: white for the zero coefficients, and red and blue for the positive and negative coefficients (with intensity proportional to the magnitude of the coefficients). The result is a “bar code”, and for the trefoil $\textcircled{3}$ it is

Similarly, θ is a 2-variable Laurent polynomial, in variables T_1 and T_2 . We can turn such a polynomial into a 2D array of coefficients and then using the same rules, into a 2D array of colours, namely, into a picture. To highlight a certain conjectured hexagonal symmetry of the resulting pictures, we apply a shear transformation to the plane before printing. So a monomial $cT_1^{n_1}T_2^{n_2}$ gets printed at position $(n_1 - n_2/2, \sqrt{3}n_2/2)$ instead of the more straightforward (n_1, n_2) . On the right is the 2D picture corresponding to the polynomial $2 + T_1 - T_1T_2 + T_2 - T_1^{-1} + T_1^{-1}T_2^{-1} - T_2^{-1}$.



Thus Θ becomes a pair of pictures: a bar code, and a 2D picture that we call a “hexagonal QR code”. For the knots in the Rolfsen table (with the unknot prepended at the start), they are in Figure 1.1. For some alternating square weave knots, they are in Figure 1.2, and for a random square weave, in Figure 1.3. In addition, the hexagonal QR codes of 15 knots with ≥ 300 crossings are in Figure 1.4, and Θ of a 132-crossing torus knot is in Figure 3.1. Some further computations and figures, also highlighting the parity of coefficients rather than just their signs, are at [Lal].

Clearly there are patterns in these figures. There is a hexagonal symmetry and the QR codes are nearly always hexagons (these are independent properties). Much more can be seen in Figure 1.1. In Figure 1.4 there seem to be large-scale patterns perhaps reminiscent of the “Chladni figures” formed by powders atop vibrating plates (on right). We can’t prove any of these things, and the last one, we can’t even formulate properly. Yet they are clearly there, too clear to be the result of chance alone.

We plan to have fun over the next few years observing and proving these patterns. We hope that others will join us too.



FIGURE 1.1. Θ as a bar code and a QR code, for all the knots in the Rolfsen table.