


CORRIGENDUM TO “FINITE TYPE INVARIANTS OF W-KNOTTED OBJECTS II”

DROR BAR-NATAN AND ZSUZSANNA DANCZO


In Section 4 of [BND] we introduce and study *w-tangled foams*. These are defined combinatorially, as a finitely generated *circuit algebra* with certain Reidemeister relations. For motivation, we present a local topological interpretation of w-tangled foams as tangled tubes in \mathbb{R}^4 . Unfortunately, some errors have occurred in this interpretation, stemming from our lack of care around “2D orientations”, as below. We wish to thank Yusuke Kuno and Haruko Miyazawa for noting these issues.

- If foams are to have 2D orientations, they cannot be glued to each other unless these orientations match. This breaks the circuit algebra structure. This can be corrected by switching to *coloured* circuit algebras, in which strands and strand-ends are coloured by their 2D orientations, and gluings are allowed only if colours match. However, we choose a different resolution explained below.
- We missed that the “ A_e ” operations of flipping only the 1D but not the 2D orientations of a strand interact unpleasantly with R4 moves, as shown on the right. This can be corrected by giving more care to the colours (2D orientations) of the edges next to a vertex and making the precise meaning of the R4 move depend on these colours. Since this is notationally tedious, we choose a different resolution below.



Although the resolution is a small change to the content of the paper, it necessitates some notation and language changes, and for the convenience of our readers we have incorporated these into the body of the paper as a revised version, available at [BNDv2].

The easier solution to both of the problems above is to forgo of 2D orientations and of the operations A_e altogether, and to replace them with more serious attention to the *wens*, which are cut open Klein-bottles ([BND, Section 4.5], and [BNDv2, Section 4.1]). Indeed, conjugating a strand by a wen and reversing it (strand reversal is the operation S_e , unchanged from the original paper) has the same effect on crossings as the operation A_e was meant to have.



The main result of the paper is identifying *homomorphic expansions* for w-tangled foams with solutions to the Kashiwara-Vergne (KV) equations. A crucial equation relating the values of such an expansion, the *unitarity equation* $V \cdot A_1 A_2(V) = 1$ [BND, Equation (12)] is replaced by an equivalent equation, $V^* V = 1$ [BNDv2, Equation (U), page 41] in which the “global adjoint” V^* is obtained from V by multiplying it by a wen on all three sides, and reversing all strand orientations.

Date: first edition October 30, 2023, this edition February 29, 2024.

In [BND] we claimed that the value of the wen under an expansion must be 1, and that expansions for w-tangled foams are in bijection with Kashiwara-Vergne solutions with *even Duflou function*. In fact, the value of the wen could be non-trivial [BNDv2, Lemma 4.9], but the statement regarding KV solutions remains true for homomorphic expansions where the value of the wen is set to be 1.

Theorem 4.9 of [BND] stated that for “orientable w-tangled foams”, i.e. where wens are not included, homomorphic expansions are in one to one correspondence with KV solutions in general. This theorem remains true: in [BNDv2, Section 4.6] we present a more careful definition of orientable w-tangled foams as a sub-circuit algebra. Edge-wise adjoint operations replaced by the well-defined global adjoint (global orientation reversal and composition with wens at every tangle end) as above, noting that these wens all cancel. Homomorphic expansions for these foams are indeed in one to one correspondence with KV solutions.

We note that the results of papers [BDS, DHR] which build on Section 4 of [BND] are unaffected by this Corrigendum: in these papers one can simply mechanically replace $A_1A_2(V)$ with V^* .

CONFLICT OF INTEREST STATEMENT

On behalf of all authors, the corresponding author states that there is no conflict of interest.

REFERENCES

- [BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects II: Tangles and the Kashiwara-Vergne Problem*, Math. Ann. **367** (2017) 1517–1586, [arXiv:1405.1955v3](https://arxiv.org/abs/1405.1955v3).
- [BNDv2] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects II: Tangles and the Kashiwara-Vergne Problem*, <http://drorbn.net/WK02> and [arXiv:1405.1955](https://arxiv.org/abs/1405.1955).
- [BDS] D. Bar-Natan, Z. Dancso, N. Scherich, *Ribbon 2-knots, $1+1=2$, and Duflou’s Theorem for Arbitrary Lie Algebras* Alg. Geom. Topol. **20**(7) 3733–3760. [arXiv:1811.08558](https://arxiv.org/abs/1811.08558)
- [DHR] Z. Dancso, I. Halacheva, M. Robertson, *A Topological Characterisation of the Kashiwara–Vergne Groups* Trans. Amer. Math. Soc. **376**(5), 3265–3317. [arXiv:2106.02373](https://arxiv.org/abs/2106.02373)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO ONTARIO M5S 2E4, CANADA
Email address: drorbn@math.toronto.edu
URL: <http://www.math.toronto.edu/~drorbn>

SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF SYDNEY, CARSLAW BUILDING,
CAMPERDOWN, NSW, 2006, AUSTRALIA
Email address: zsuzsanna.dancso@sydney.edu.au