

Greece handout on July 15, 2016 (as printed)

July 17, 2016 8:05 PM

Dror Bar-Natan: Talks: Greece-1607: <http://drorbn.net/Greece-1607/>
 Work in Progress! **The Brute and the Hidden Paradise**

Abstract. There is expected to be a hidden paradise of poly-time computable knot polynomials lying just beyond the Alexander polynomial. I will describe my brute attempts to gain entry.

Why "expected"? Gauss diagram $v_{d,f}(K) = \sum_{Y \subset X(K), |Y|=d} f(Y)$ formulas [PV, GPV] show that finite-type invariants are all poly-time, and tempt to conjecture that there are no others. But Alexander shows it nonsense:

d	2	3	4	5	6	7	8	...
known invts* in $O(n^d)$	1	1	∞	3	4	8	11	...

This is an unreasonable picture! *Fresh, numerical, no cheating. So there ought to be further poly-time invariants.


Also. • The line above the Alexander line in the Melvin-Morton [MM, Ro] expansion of the coloured Jones polynomial. • The 2-loop contribution to the Kontsevich integral.

Why "paradise"? Foremost answer: **OBVIOUSLY.** Cf. proving (incomputable A)=(incomputable B), or categorifying (incomputable C).

$\omega\beta/K17$:

(extend to tangles, perhaps detect non-slice ribbon knots)

Moral. Need "stitching":



Definition 1. A "ribbon knot" is a knot that can be presented in the boundary of a disk D^2 which is allowed to have "ribbon singularities" but not "sharp singularities". See Figure 2.

Figure 2. A ribbon singularity is a sharp singularity, and an example of a ribbon knot.

Definition 2. Let \mathcal{T}_2 denote the set of all tangles T with $2n$ components that connect $2n$ points along a "top end" with $2n$ points along a "bottom end" inducing the identity permutation of ends (an example is the tangle in Figure 1). Given $T \in \mathcal{T}_2$, let $\sigma(T)$ be the result of stitching its components at the top in pairs as on the right - it is an n -component tangle all of whose ends are at the bottom, and we (conveniently) denote the set of all such by \mathcal{T}_n , so $\sigma: \mathcal{T}_2 \rightarrow \mathcal{T}_n$. Likewise let $\sigma(T)$ be the result of stitching T both at the top and at the bottom, also as on the right. So $\sigma(T)$ is a 1-component tangle, which is the same as a knot, and $\sigma: \mathcal{T}_2 \rightarrow \mathcal{T}_1$.

Theorem 1 (I have not seen this theorem in the literature, yet it is not difficult to prove). The set of ribbon knots is the set of all knots K that can be written as $K = \sigma(T)$ for which $\sigma(T)$ is the unknotted (non-singular) tangle $\sigma(T)$.

$(\text{ribbon knots}) = \{ \sigma(T) : T \in \mathcal{T}_2, \text{ and } \sigma(T) \text{ is unknotted} \}$

Now suppose we have an invariant $Z: \mathcal{T}_1 \rightarrow A_1$ of tangles, which takes values in some spaces A_i . Suppose also we have operators $\tau_1: A_2 \rightarrow A_1$ and $\tau_2: A_3 \rightarrow A_1$ such that the diagram on the right is commutative. Then

$$Z(\text{ribbon knot}) \in \mathcal{R}_1 := \{ \tau_1(Z) : Z \in A_2, \text{ and } \tau_2(Z) \in A_1 \} \subset A_1 \quad (1)$$

$$\mathcal{R}_2 := \{ \tau_1(Z) : Z \in A_3, \text{ and } \tau_2(Z) \in A_1 \} \subset A_1 \quad (2)$$

where $\tau_1 = ZU \in A_1$. If the target spaces A_i are algebraic (polynomials, matrices, matrices of polynomials, etc.) and the operators τ_1 and τ_2 are algebraic maps between them (at this stage, meaning just "nice simple algebraic formulae"), then \mathcal{R}_1 is an algebraically defined set. Hence we potentially have an algebraic way to detect non-ribbon knots: if $Z(K) \notin \mathcal{R}_1$, then K is not ribbon.

As it turns out, it is valuable to detect non-ribbon knots. Indeed the Slice-Ribbon Conjecture (Fox, 1980b) asserts that every slice knot is (at least in S^3) that can be presented as the boundary of a disk embedded in \mathbb{R}^4 is ribbon. Gompf, Schleichmann, and Thompson (GST) describe a family of slice knots which they conjecture are not ribbon (the simplest of these is on the right). With the algebraic technology described above it may be possible to show that the [GST] knots are indeed non-ribbon, thus disproving the Slice-Ribbon Conjecture.

What would it take?

- C1. An invariant Z which makes sense on tangles and for which diagram (1) commutes.
- C2. Z cannot be a simple extension of the Alexander polynomial to tangles, for by Fox-Milnor [FM] the Alexander polynomial does not detect non-ribbon slice knots.
- C3. Z cannot be computable from finitely many finite type invariants, for this would contradict the results of Ng [Ng].
- C4. Z must be computable on at least the simplest [GST] knot, which has 48 crossings.
- C5. It is better if in some meaningful sense the size of the space A_i grows slowly in i . Indeed in C2, if A_2 is much bigger than A_1 and A_3 then at least generally \mathcal{R}_1 will be the full set A_1 and our condition will be empty.

No invariant that I know now meets these criteria. Alexander and Vassiliev fail C2 and C3, respectively. Almost all quantum invariants and link homomorphisms pass C1-C3, but fail C4. HOMFLY-PT and Khovanov potentially pass C4, yet fail C5. We must come up with something new.

[EM] R. E. Fox and J. W. Milnor. Singularities of 2-Spheres in 4-Space and Calculation of Knots. Osaka J. Math. 3 (1966) 257-267.
 [GST] R. E. Gompf, M. Schleichmann, and A. Thompson. *Filtered Knots and Potential Counterexamples to the Property 2R and Slice-Ribbon Conjectures*. Geom. and Top. 14 (2010) 2305-2347. arXiv:1005.1601.
 [Ng] K. Y. Ng. *Groups of ribbon knots*. Topology 37 (1998) 441-458. arXiv:alg-geom/9802017 (with an addendum at arXiv:math.GT/01010074)

*A slight subtlety above: There is no taking limits here, and C1 does not preclude the possibility that Z is computable from infinitely many finite type invariants. The first Milnor condition on the Alexander polynomial of ribbon knots, for example, is expressible in terms of the full Alexander polynomial, yet not in terms of any finite number thereof. Unfortunately by C2 it cannot be used here.

Why "brute"? Cause it's the only thing I know, for now. There may be better ways in, and it's fair to hope that sooner or later they will be found.

The Gold Standard is set by the formulas [BNS, BN] for Alexander. An S -component tangle T has $\Gamma(T) \in R_S \times M_S \times S(R_S) = \left\{ \frac{\omega}{S} \frac{S}{A} \right\}$ with $R_S := \mathbb{Z}\langle t_a : a \in S \rangle$:

$$\left(\begin{array}{c} a \\ \nearrow \\ b \end{array} \otimes \begin{array}{c} b \\ \nearrow \\ a \end{array} \right) \rightarrow \begin{array}{c|c} 1 & a \\ \hline a & 1 - t_a^{-1} \\ \hline b & 0 \\ & t_a^{-1} \end{array} \quad T_1 \sqcup T_2 \rightarrow \begin{array}{c|c} \omega_1 \omega_2 & S_1 \\ \hline S_1 & A_1 \\ \hline S_2 & 0 \\ & A_2 \end{array}$$


$$\begin{array}{c|c} \omega & a \quad b \quad S \\ \hline a & \alpha \quad \beta \quad \theta \\ b & \gamma \quad \delta \quad \epsilon \\ S & \phi \quad \psi \quad \Xi \end{array} \xrightarrow{m_c^{ab}} \begin{array}{c|c} (1-\beta)\omega & c \quad S \\ \hline c & \gamma + \frac{\alpha\delta}{1-\beta} \quad \epsilon + \frac{\delta\theta}{1-\beta} \\ S & \phi + \frac{\alpha\psi}{1-\beta} \quad \Xi + \frac{\psi\theta}{1-\beta} \end{array}$$

$t_a, t_b \rightarrow t_c$

Help Needed!
 I'm slow and feeble-minded.

For long knots, ω is Alexander, and that's the fastest Alexander algorithm I know!
 Dunfield: 1000-crossing fast.

Theorem [EK, Ha, En, Se]. There is a "homomorphic expansion"




(it is enough to know Z on \mathcal{A}_S^v and have disjoint union and stitching formulas) ... exponential and too hard!

Idea. Look for "ideal" quotients of \mathcal{A}_S^v that have poly-sized descriptions; ... specifically, limit the co-brackets.

1-co and 2-co, aka TC and TC², on the right. The primitives that remain are:




The 2D relations come from the relation with 2D Lie bialgebras:



We let $\mathcal{A}^{2,2}$ be \mathcal{A}^v modulo 2-co and 2D, and $\mathcal{P}^{2,2}$ be the projection of log Z to $\mathcal{P}^{2,2} := \pi \mathcal{P}^v$, where \mathcal{P}^v are the primitives of \mathcal{A}^v .

Main Claim. $\mathcal{P}^{2,2}$ is poly-time computable.

Main Point. $\mathcal{P}^{2,2}$ is poly-size, so how hard can it be? Indeed, as a module over $\mathbb{Q}\langle b_i \rangle$, $\mathcal{P}^{2,2}$ is at most




Claim. $R_{jk} = e^{a_{jk}} e^{\rho_{jk}}$ is a solution of the Yang-Baxter / R3 equation $R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}$ in $\exp \mathcal{P}^{2,2}$, with $\rho_{jk} :=$

$$\psi(b_j) \left(-c_k + \frac{c_k a_{jk}}{b_j} - \frac{\delta a_{jk} a_{jk}}{b_j^2} \right) + \frac{\phi(b_j) \psi(b_k)}{b_k \phi(b_k)} \left(c_k a_{kk} - \frac{\delta a_{jk} a_{kk}}{b_j} \right),$$

and with $\phi(x) := e^{-x} - 1 = -x + x^2/2 - \dots$, and $\psi(x) := ((x+2)e^{-x} - 2 + x)/(2x) = x^2/12 - x^3/24 + \dots$. (This already gives some new (v)-braid group representations, as below).

Problem. How do we multiply in $\exp(\mathcal{P}^{2,2})$? How do we stitch? BCH is a theoretical dream. Instead, use "scatter and glow" and "feedback loops":



The Brute and the Hidden Paradise

Local Algebra (with van der Veen) Much can be re-formulated as (non-standard) “quantum algebra” for the 4D Lie algebra $\mathfrak{g} = \langle b, c, u, w \rangle$ over $\mathbb{Q}[\epsilon]/(\epsilon^2 = 0)$, with b central and $[w, c] = w$, $[c, u] = u$, and $[u, w] = b - 2\epsilon c$. The key: $a_{ij} = (b_i - \epsilon c_i)c_j + u_i w_j$ in $\mathcal{U}(\mathfrak{g})^{\otimes(i,j)}$.



van der Veen

Some (new) representations of the (v-)braid groups. oeβ/Reps Bureau (old)

```
B_{i,j}[\xi] := \xi / . v_j \mapsto (1-t) v_i + t v_j
Column@{lhs = {v1, v2, v3} // B_{1,2} // B_{1,3} // B_{2,3},
  rhs = {v1, v2, v3} // B_{2,3} // B_{1,3} // B_{1,2},
  lhs - rhs // Expand}
```

```
{v1, (1-t) v1 + t v2, (1-t) v1 + t ((1-t) v2 + t v3)}
{v1, (1-t) v1 + t v2,
 (1-t) ((1-t) v1 + t v2) + t ((1-t) v1 + t v3)}
{0, 0, 0}
```

$G_{i,j}[\xi] := \xi / . v_j \mapsto (1-t_i) v_i + t_i v_j$ Gassner (old) ... Overcrossings Commute (OC)

```
Column@{lhs = {v1, v2, v3} // G_{1,2} // G_{1,3},
  Expand[lhs - ({v1, v2, v3} // G_{1,3} // G_{1,2})]}
{v1, (1-t1) v1 + t1 v2, (1-t1) v1 + t1 v3}
{0, 0, 0}
```

... Undercrossings Commute (UC):

```
Column@{lhs = {v1, v2, v3} // G_{1,3} // G_{2,3},
  rhs = {v1, v2, v3} // G_{2,3} // G_{1,3},
  lhs - rhs // Expand}
{v1, v2, (1-t1) v1 + t1 ((1-t2) v2 + t2 v3)}
{v1, v2, (1-t2) v2 + t2 ((1-t1) v1 + t1 v3)}
{0, 0, v1 - t1 v1 - t2 v1 + t1 t2 v1 - v2 + t1 v2 + t2 v2 - t1 t2 v2}
```

Gassner Plus (new?)

```
GP_{i,j}[\xi] := Expand[\xi / . {u_j \mapsto (1-t_i) u_i + t_i u_j,
  f . v_j \mapsto f (1-t_i) v_i + f t_i v_j + (t_i - 1) (t_i \partial_{c_i} f - t_j \partial_{c_j} f) u_i +
  f t_i u_i}];
```

```
bas = {f[t1, t2, t3] v1, f[t1, t2, t3] v2, f[t1, t2, t3] v3,
  u1, u2, u3};
Short[lhs = bas // GP_{1,2} // GP_{1,3} // GP_{2,3}, 2] ... R3 (left)
```

```
{f[t1, t2, t3] v1, f[t1, t2, t3] t1 u1 + f[t1, t2, t3] v1 -
  f[t1, t2, t3] t1 v1 + <<6>> + t1^2 u1 f^{(1,0,0)}[t1, t2, t3],
 <<1>> + <<19>> + <<1>>, <<1>>, u1 - t1 u1 + t1 u2,
 u1 - t1 u1 + t1 u2 - t1 t2 u2 + t1 t2 u3}
```

```
(bas // GP_{2,3} // GP_{1,3} // GP_{1,2}) - lhs ... R3 (rest)
{0, 0, 0, 0, 0, 0}
(bas // GP_{1,2} // GP_{1,3}) - (bas // GP_{1,3} // GP_{1,2}) ... OC
{0, 0, 0, 0, 0, 0}
```

Question. Does Gassner Plus factor through Gassner?

$K\delta_{i,j} := \text{KroneckerDelta}[i, j];$ Turbo-Gassner (new!)

```
TG_{i,j}[\xi] := Expand[\xi / . {
  f . v_k \mapsto Plus[f v_k / . v_j \mapsto (1-t_i) v_i + t_i v_j,
  (1-t_i^{-1}) (t_i \partial_{c_i} f - t_j \partial_{c_j} f) *
  (u_k / . u_j \mapsto (1-t_i) u_i + t_i u_j) * u_i w_j,
  K\delta_{k,i} f (u_j - u_i) u_i w_j,
  u_j \mapsto (1-t_i) u_i + t_i u_j,
  w_j \mapsto w_i + (1-t_i^{-1}) w_j, w_j \mapsto t_i^{-1} w_j}];
```

```
bas = {f[t1, t2, t3] v1, f[t1, t2, t3] v2, f[t1, t2, t3] v3,
  u1, u2, u3, w1, w2, w3};
```

```
(bas // TG_{1,2} // TG_{1,3} // TG_{2,3}) - (bas // TG_{2,3} // TG_{1,3} // TG_{1,2}) . R3
{0, 0, 0, 0, 0, 0, 0, 0}
(bas // TG_{1,2} // TG_{1,3}) - (bas // TG_{1,3} // TG_{1,2}) ... OC
```

```
{0, -f[t1, t2, t3] u1 u2 w3 + f[t1, t2, t3] t1 u1 u2 w3 +
  f[t1, t2, t3] u1 u3 w3 - f[t1, t2, t3] t1 u1 u3 w3,
 -f[t1, t2, t3] u1 u2 w2 + f[t1, t2, t3] t1 u1 u2 w2 +
  f[t1, t2, t3] u1 u3 w2 -
  f[t1, t2, t3] t1 u1 u3 w2, 0, 0, 0, 0, 0, 0}
```

$\eta / : \eta[i_-]^2 = 0; \eta / : \eta[i_-] \eta[j_-] = 0;$ Turbo-Bureau (new!)

```
TB_{i,j}[\xi] :=
Expand[\xi / . {
  f . v_k \mapsto Plus[f v_k / . v_j \mapsto (1-t-\eta[i]) v_i + (t+\eta[i]) v_j,
  (t-1) (Coefficient[f, \eta[i]] - Coefficient[f, \eta[j]]) *
  (u_k / . u_j \mapsto (1-t) u_i + t u_j) * u_i w_j,
  K\delta_{k,i} (f / . \eta \rightarrow 0) (u_j - u_i) u_i w_j,
  u_j \mapsto (1-t) u_i + t u_j,
  w_i \mapsto w_i + (1-t^{-1}) w_j, w_j \mapsto t^{-1} w_j}];
```

```
ff = f0 + f1 \eta[1] + f2 \eta[2] + f3 \eta[3];
bas = {ff v1, ff v2, ff v3, u1^2 w1, u2^2 w2, u1, u2, u3, w1, w2, w3};
Short[lhs = bas // TB_{1,2} // TB_{1,3}, 3] ... OC
```

```
{f0 v1 - f0 u1^2 w2 - f1 u1^2 w2 + t f1 u1^2 w2 + f2 u1^2 w2 - t f2 u1^2 w2 +
  f0 u1 u2 w2 - f0 u1^2 w3 - f1 u1^2 w3 + t f1 u1^2 w3 + f3 u1^2 w3 - t f3 u1^2 w3 +
  f0 u1 u3 w3 + f1 v1 \eta[1] + f2 v1 \eta[2] + f3 v1 \eta[3], <<9>>, \frac{w3}{t}}
```

```
rhs = bas // TB_{1,3} // TB_{1,2}; lhs - rhs
{0, -f0 u1 u2 w3 + t f0 u1 u2 w3 + f0 u1 u3 w3 - t f0 u1 u3 w3,
 -f0 u1 u2 w2 + t f0 u1 u2 w2 + f0 u1 u3 w2 - t f0 u1 u3 w2,
 0, 0, 0, 0, 0, 0, 0}
```

References.

[BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, oeβ/KBH, arXiv: 1308.1721.

[BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I, II, IV*, oeβ/WKO1, oeβ/WKO2, oeβ/WKO4, arXiv:1405.1956, arXiv: 1405.1955, arXiv:1511.05624.

[BNG] D. Bar-Natan and S. Garoufalidis, *On the Melvin-Morton-Rozansky conjecture*, Invent. Math. **125** (1996) 103–133.

[BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.

[En] B. Enriquez, *A Cohomological Construction of Quantization Functors of Lie Bialgebras*, Adv. in Math. **197-2** (2005) 430-479, arXiv:math/0212325.

[EK] P. Etingof and D. Kazhdan, *Quantization of Lie Bialgebras, I*, Selecta Mathematica **2** (1996) 1–41, arXiv:q-alg/9506005.

[GPV] M. Goussarov, M. Polyak, and O. Viro, *Finite type invariants of classical and virtual knots*, Topology **39** (2000) 1045–1068, arXiv: math.GT/9810073.

[Ha] A. Haviv, *Towards a diagrammatic analogue of the Reshetikhin-Turaev link invariants*, Hebrew University PhD thesis, Sep. 2002, arXiv: math.QA/0211031.

[MM] P. M. Melvin and H. R. Morton, *The coloured Jones function*, Commun. Math. Phys. **169** (1995) 501–520.

[PV] M. Polyak and O. Viro, *Gauss Diagram Formulas for Vassiliev Invariants*, Inter. Math. Res. Notices **11** (1994) 445–453.

[Ro] L. Rozansky, *A contribution of the trivial flat connection to the Jones polynomial and Witten's invariant of 3d manifolds, I*, Comm. Math. Phys. **175-2** (1996) 275–296, arXiv:hep-th/9401061.

[Se] P. Ševera, *Quantization of Lie Bialgebras Revisited*, Sel. Math., NS, to appear, arXiv:1401.6164.



“God created the knots, all else in topology is the work of mortals.”

Leopold Kronecker (modified)



1. Roland's typo.
2. Picture from PolyPoly meetings.
3. Euler comment: $Ef := (\deg f)f$, $Ee^x = xe^x$, $E(e^x e^{ye^z}) = xe^x e^{ye^z} + e^x ye^z + \dots$
4. Some flowers?