

**Cars, Interchanges, Traffic Counters, and some Pretty Darned Good Knot Invariants** More at $\omega\epsilon\beta/\text{APAI}$


Abstract. Reporting on joint work with Roland van der Veen, I'll tell you some stories about ρ_1 , an easy to define, strong, fast to compute, homomorphic, and well-connected knot invariant. ρ_1 was first studied by Rozansky and Overbay [Ro1, Ro2, Ro3, Ov] and Ohtsuki [Oh2], it has far-reaching generalizations, it is elementary and dominated by the coloured Jones polynomial, and I wish I understood it.

Common misconception. Dominated, elementary $\Rightarrow$ lesser.


We seek strong, fast, homomorphic knot and tangle invariants.

Strong. Having a small “kernel”.


Fast. Computable even for large knots (best: poly time).



Homomorphic. Extends to tangles and behaves under tangle operations; especially gluings and doublings:



Why care for “Homomorphic”? **Theorem.** A knot K is ribbon iff there exists a $2n$ -component tangle T with skeleton as below such that $\tau(T) = K$ and where $\delta(T) = U$ is the untangle:



Hear more at $\omega\epsilon\beta/\text{AKT}$.

Acknowledgement. This work was supported by NSERC grant RGPIN-2018-04350 and by the Chu Family Foundation (NYC).

[BV1] D. Bar-Natan and R. van der Veen, *A Polynomial Time Knot Polynomial*, Proc. Amer. Math. Soc. **147** (2019) 377–397, [arXiv:1708.04853](https://arxiv.org/abs/1708.04853).

[BV2] D. Bar-Natan and R. van der Veen, *Perturbed Gaussian Generating Functions for Universal Knot Invariants*, [arXiv:2109.02057](https://arxiv.org/abs/2109.02057).

[Dr] V. G. Drinfel'd, *Quantum Groups*, Proc. Int. Cong. Math., 798–820, Berkeley, 1986.

[Jo] V. F. R. Jones, *Hecke Algebra Representations of Braid Groups and Link Polynomials*, Annals Math., **126** (1987) 335–388.

[La] R. J. Lawrence, *Universal Link Invariants using Quantum Groups*, Proc. XVII Int. Conf. on Diff. Geom. Methods in Theor. Phys., Chester, England, August 1988. World Scientific (1989) 55–63.

[LTW] X.-S. Lin, F. Tian, and Z. Wang, *Burau Representation and Random Walk on String Links*, Pac. J. Math., **182-2** (1998) 289–302, [arXiv:q-alg/9605023](https://arxiv.org/abs/q-alg/9605023).

[Oh1] T. Ohtsuki, *Quantum Invariants*, Series on Knots and Everything **29**, World Scientific 2002.

[Oh2] T. Ohtsuki, *On the 2-loop Polynomial of Knots*, Geom. Top. **11** (2007) 1357–1475.

[Ov] A. Overbay, *Perturbative Expansion of the Colored Jones Polynomial*, Ph.D. thesis, University of North Carolina, August 2013, $\omega\epsilon\beta/\text{Ov}$.

[Ro1] L. Rozansky, *A Contribution of the Trivial Flat Connection to the Jones Polynomial and Witten's Invariant of 3D Manifolds, I*, Comm. Math. Phys. **175-2** (1996) 275–296, [arXiv:hep-th/9401061](https://arxiv.org/abs/hep-th/9401061).

[Ro2] L. Rozansky, *The Universal R-Matrix, Burau Representation and the Melvin-Morton Expansion of the Colored Jones Polynomial*, Adv. Math. **134-1** (1998) 1–31, [arXiv:q-alg/9604005](https://arxiv.org/abs/q-alg/9604005).

[Ro3] L. Rozansky, *A Universal $U(1)$ -RCC Invariant of Links and Rationality Conjecture*, [arXiv:math/0201139](https://arxiv.org/abs/math/0201139).

[Sch] S. Schaveling, *Expansions of Quantum Group Invariants*, Ph.D. thesis, Universiteit Leiden, September 2020, $\omega\epsilon\beta/\text{Scha}$.


* In algebra $x \sim 0$ if for every y in the ideal generated by x , $1 - y$ is invertible.



Jones:

Formulas stay; interpretations change with time.

Formulas. Draw an n -crossing knot K as on the right: all crossings face up, and the edges are marked with a running index $k \in \{1, \dots, 2n + 1\}$ and with rotation numbers φ_k . Let A be the $(2n + 1) \times (2n + 1)$ matrix constructed by starting with the identity matrix I , and adding a 2×2 block for each crossing:



$$c: \begin{array}{c} s = +1 \\ j+1 \uparrow \quad i+1 \uparrow \\ i \quad j \end{array} \rightarrow \begin{array}{c} s = -1 \\ i+1 \uparrow \quad j+1 \uparrow \\ j \quad i \end{array} \rightarrow \begin{array}{c|cc} A & \text{col } i+1 & \text{col } j+1 \\ \hline \text{row } i & -T^s & T^s - 1 \\ \text{row } j & 0 & -1 \end{array}$$

Let $G = (g_{\alpha\beta}) = A^{-1}$. For the trefoil example, it is:

$$A = \begin{pmatrix} 1 & -T & 0 & 0 & T-1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -T & 0 & 0 & T-1 \\ 0 & 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & T-1 & 0 & 1 & -T & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$G = \begin{pmatrix} 1 & T & 1 & T & 1 & T & 1 \\ 0 & 1 & \frac{1}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T^2}{T^2-T+1} & 1 \\ 0 & 0 & \frac{T^2-T+1}{1-T} & \frac{T}{T^2-T+1} & \frac{T^2-T+1}{T} & \frac{T^2-T+1}{T} & 1 \\ 0 & 0 & \frac{T^2-T+1}{1-T} & \frac{T^2-T+1}{T} & \frac{T^2-T+1}{T} & \frac{T^2-T+1}{T} & 1 \\ 0 & 0 & \frac{1-T}{T^2-T+1} & -\frac{(T-1)T}{T^2-T+1} & \frac{1}{T^2-T+1} & \frac{T}{T^2-T+1} & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

“The Green Function”

Note. The Alexander polynomial Δ is given by

$$\Delta = T^{(-\varphi-w)/2} \det(A), \quad \text{with } \varphi = \sum_k \varphi_k, \quad w = \sum_c s.$$

Classical Topologists: This is boring. Yawn.

Formulas, continued. Finally, set

$$R_1(c) := s(g_{ji}(g_{j+1,j} + g_{j,j+1} - g_{ij}) - g_{ii}(g_{j,j+1} - 1) - 1/2)$$

$$\rho_1 := \Delta^2 \left(\sum_c R_1(c) - \sum_k \varphi_k (g_{kk} - 1/2) \right).$$

In our example $\rho_1 = -T^2 + 2T - 2 + 2T^{-1} - T^{-2}$.

Theorem. ρ_1 is a knot invariant.

Proof: later.

Classical Topologists: Whiskey Tango Foxtrot?

Cars, Interchanges, and Traffic Counters.

Cars always drive forward. When a car crosses over a bridge it goes through with (algebraic) probability $T^s \sim 1$, but falls off with probability $1 - T^s \sim 0^*$. At the very end, cars fall off and disappear. See also [Jo, LTW].




image credits: diamondtraffic.com

image credits: Dall-E

Preliminaries

This is Rho.nb of <http://drorbn.net/oa22/ap>.

Once[<< KnotTheory` ; << Rot.m];

Loading KnotTheory` version

of February 2, 2020, 10:53:45.2097.

Read more at <http://katlas.org/wiki/KnotTheory>.

Loading Rot.m from <http://drorbn.net/la22/ap>

to compute rotation numbers.

The Program

```
R1[s_, i_, j_] :=
  S (gji (gj+,j + gj,j+ - gij) - gi (gj,j+ - 1) - 1/2);
Z[K_] := Module[{Cs, φ, n, A, s, i, j, k, Δ, G, ρ1},
  {Cs, φ} = Rot[K]; n = Length[Cs];
  A = IdentityMatrix[2 n + 1];
  Cases[Cs, {s_, i_, j_} >=
    (A[[{i, j}, {i + 1, j + 1}]] += ( -T^s T^s - 1 ))];
  Δ = T^(-Total[φ] - Total[Cs[[All, 1]]]) / 2 Det[A];
  G = Inverse[A];
  ρ1 = Sum_{k=1}^n R1 @@ Cs[[k]] - Sum_{k=1}^{2 n} φ[[k]] (gkk - 1/2);
  Factor@
    {Δ, Δ^2 ρ1 /. α_+ >= α + 1 /. g_{α,β} >= G[[α, β]]};
```

The First Few Knots


```
TableForm[Table[Join[{K[[1]][[k][2]], Z[K]},
  {K, AllKnots[{3, 6}]}], TableAlignments -> Center]
```

3 ₁	$\frac{1-T+T^2}{T}$	$\frac{(-1+T)^2 (1+T^2)}{T^2}$
4 ₁	$-\frac{1-3T+T^2}{T}$	0
5 ₁	$\frac{1-T+T^2-T^3+T^4}{T^2}$	$\frac{(-1+T)^2 (1+T^2) (2+T^2+2T^4)}{T^4}$
5 ₂	$\frac{2-3T+2T^2}{T}$	$\frac{(-1+T)^2 (5-4T+5T^2)}{T^2}$
6 ₁	$-\frac{(-2+T) (-1+2T)}{T}$	$\frac{(-1+T)^2 (1-4T+T^2)}{T^2}$
6 ₂	$-\frac{1-3T+3T^2-3T^3+T^4}{T^2}$	$\frac{(-1+T)^2 (1-4T+4T^2-4T^3+4T^4-4T^5+T^6)}{T^4}$
6 ₃	$\frac{1-3T+5T^2-3T^3+T^4}{T^2}$	0



$$p = 1 - T^s$$

Fast!



Timing@

```
Z[GST48 = EPD[X14,1, X2,29, X3,40, X43,4, X26,5, X6,95,
  X96,7, X13,8, X9,28, X10,41, X42,11, X27,12, X30,15,
  X16,61, X17,72, X18,83, X19,34, X89,20, X21,92,
  X79,22, X68,23, X57,24, X25,56, X62,31, X73,32,
  X84,33, X50,35, X36,81, X37,70, X38,59, X39,54, X44,55,
  X58,45, X69,46, X80,47, X48,91, X90,49, X51,82, X52,71,
  X53,60, X63,74, X64,85, X76,65, X87,66, X67,94,
  X75,86, X88,77, X78,93]]
```

$$\{170.313, \left\{ -\frac{1}{T^8} (-1 + 2T - T^2 - T^3 + 2T^4 - T^5 + T^8) \right.$$

$$\left. (-1 + T^3 - 2T^4 + T^5 + T^6 - 2T^7 + T^8) \right\}, \frac{1}{T^{16}}$$

$$(-1 + T)^2 (5 - 18T + 33T^2 - 32T^3 + 2T^4 + 42T^5 - 62T^6 - 8T^7 + 166T^8 - 242T^9 + 108T^{10} + 132T^{11} - 226T^{12} + 148T^{13} - 11T^{14} - 36T^{15} - 11T^{16} + 148T^{17} - 226T^{18} + 132T^{19} + 108T^{20} - 242T^{21} + 166T^{22} - 8T^{23} - 62T^{24} + 42T^{25} + 2T^{26} - 32T^{27} + 33T^{28} - 18T^{29} + 5T^{30}) \}$$

Strong!

```
{NumberOfKnots[{3, 12}],
```

```
Length@
```

```
Union@Table[Z[K], {K, AllKnots[{3, 12}]}],
```

```
Length@
```

```
Union@Table[{HOMFLYPT[K], Kh[K]},
  {K, AllKnots[{3, 12}]}]]
```

```
{2977, 2882, 2785}
```

So the pair (Δ, ρ_1) attains 2,882 distinct values on the 2,977 prime knots with up to 12 crossings (a deficit of 95), whereas the pair (HOMFLYPT, Khovanov Homology) attains only 2,785 distinct values on the same knots (a deficit of 192).



Hoste

Ocneanu

Millett

Freyd

Lickorish


Yetter

Przytycki

Traczyk

Khovanov

Theorem. The Green function $g_{\alpha\beta}$ is the reading of a traffic counter at β , if car traffic is injected at α (if $\alpha = \beta$, the counter is *after* the injection point).



Example.

$$\sum_{p \geq 0} (1-T)^p = T^{-1} \quad T^{-1} \quad 0 \quad 1 \quad 0 \quad 1 \quad 0 \quad 1 \quad G = \begin{pmatrix} 1 & T^{-1} & 1 \\ 0 & T^{-1} & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

Proof. Near a crossing c with sign s , incoming upper edge i and incoming lower edge j , both sides satisfy the g -rules:




$g_{i\beta} = \delta_{i\beta} + T^s g_{i+1,\beta} + (1-T^s) g_{j+1,\beta}$, $g_{j\beta} = \delta_{j\beta} + g_{j+1,\beta}$, and always, $g_{\alpha,2n+1} = 1$: use common sense and $AG = I (= GA)$.

Bonus. Near c , both sides satisfy the further g -rules:

$$g_{\alpha i} = T^{-s}(g_{\alpha,i+1} - \delta_{\alpha,i+1}), \quad g_{\alpha j} = g_{\alpha,j+1} - (1-T^s)g_{\alpha i} - \delta_{\alpha,j+1}.$$

Invariance of ρ_1 . We start with the hardest, Reidemeister 3:



⇒ Overall traffic patterns are unaffected by Reid3!

⇒ Green's $g_{\alpha\beta}$ is unchanged by Reid3, provided the cars injection site α and the traffic counters β are away.

⇒ Only the contribution from the R_1 terms within the Reid3 move matters, and using g -rules the relevant $g_{\alpha\beta}$'s can be pushed outside of the Reid3 area:

$$\delta_{i,j} := \text{If}[i == j, 1, 0];$$

$$\text{gRules}_{s,i,j} :=$$

$$\{g_{i\beta} \mapsto \delta_{i\beta} + T^s g_{i+1,\beta} + (1-T^s) g_{j+1,\beta}, g_{j\beta} \mapsto \delta_{j\beta} + g_{j+1,\beta}, g_{\alpha,i} \mapsto T^{-s}(g_{\alpha,i+1} - \delta_{\alpha,i+1}), g_{\alpha,j} \mapsto g_{\alpha,j+1} - (1-T^s)g_{\alpha i} - \delta_{\alpha,j+1}\}$$

$$\text{lhs} = R_1[1, j, k] + R_1[1, i, k^+] + R_1[1, i^+, j^+] //.$$

$$\text{gRules}_{1,j,k} \cup \text{gRules}_{1,i,k^+} \cup \text{gRules}_{1,i^+,j^+};$$

$$\text{rhs} = R_1[1, i, j] + R_1[1, i^+, k] + R_1[1, j^+, k^+] //.$$

$$\text{gRules}_{1,i,j} \cup \text{gRules}_{1,i^+,k} \cup \text{gRules}_{1,j^+,k^+};$$

Simplify[lhs == rhs]

True


Next comes Reid1, where we use results from an earlier example:

$$R_1[1, 2, 1] - 1 (g_{22} - 1/2) /. g_{\alpha,\beta} \mapsto \begin{pmatrix} 1 & T^{-1} & 1 \\ 0 & T^{-1} & 1 \\ 0 & 0 & 1 \end{pmatrix} [[\alpha, \beta]]$$

$$\frac{1}{T^2} - \frac{1}{T} - \frac{-1 + \frac{1}{T}}{T} = 0$$

Invariance under the other moves is proven similarly.

Wearing my Topology hat the formula for R_1 , and even the idea to look for R_1 , remain a complete mystery to me.



Wearing my Quantum Algebra hat, I spy a Heisenberg algebra $\mathbb{H} = A\langle p, x \rangle / ([p, x] = 1)$:

$$\text{cars} \leftrightarrow p \quad \text{traffic counters} \leftrightarrow x$$

Where did it come from? Consider $g_\epsilon := s_{2+}^\epsilon := L\langle y, b, a, x \rangle$ with relations

$$[b, x] = \epsilon x, \quad [b, y] = -\epsilon y, \quad [b, a] = 0,$$

$$[a, x] = x, \quad [a, y] = -y, \quad [x, y] = b + \epsilon a.$$

At invertible ϵ , it is isomorphic to sl_2 plus a central factor, and it can be quantized à la Drinfel'd [Dr] much like sl_2 to get an algebra $QU = A\langle y, b, a, x \rangle$ subject to (with $q = e^{\hbar\epsilon}$):

$$[b, a] = 0, \quad [b, x] = \epsilon x, \quad [b, y] = -\epsilon y,$$

$$[a, x] = x, \quad [a, y] = -y, \quad xy - qyx = \frac{1 - e^{-\hbar(b+\epsilon a)}}{\hbar}.$$

Now QU has an R -matrix solving Yang-Baxter (meaning Reid3),

$$R = \sum_{m,n \geq 0} \frac{y^m b^m \otimes (\hbar a)^m (\hbar x)^n}{m! [n]_q!}, \quad ([n]_q! \text{ is a "quantum factorial"})$$

and so it has an associated "universal quantum invariant" à la Lawrence and Ohtsuki [La, Oh1], $Z_\epsilon(K) \in QU$.

Now $QU \cong \mathcal{U}(g_\epsilon)$ (only as algebras!) and $\mathcal{U}(g_\epsilon)$ represents into $\mathbb{H}$ via

$$y \rightarrow -tp - \epsilon \cdot xp^2, \quad b \rightarrow t + \epsilon \cdot xp, \quad a \rightarrow xp, \quad x \rightarrow x,$$

(abstractly, g_ϵ acts on its Verma module

$$\mathcal{U}(g_\epsilon) / (\mathcal{U}(g_\epsilon)\langle y, a, b - \epsilon a - t \rangle) \cong \mathbb{Q}[x]$$

by differential operators, namely via $\mathbb{H}$), so R can be pushed to $\mathcal{R} \in \mathbb{H} \otimes \mathbb{H}$.

Everything still makes sense at $\epsilon = 0$ and can be expanded near $\epsilon = 0$ resulting with $\mathcal{R} = \mathcal{R}_0(1 + \epsilon \mathcal{R}_1 + \dots)$, with $\mathcal{R}_0 = e^{t(xp \otimes 1 - x \otimes p)}$ and $\mathcal{R}_1$ a quartic polynomial in p and x . So p 's and x 's get created along K and need to be pushed around to a standard location ("normal ordering"). This is done using

$$(p \otimes 1)\mathcal{R}_0 = \mathcal{R}_0(T(p \otimes 1) + (1 - T)(1 \otimes p)),$$

$$(1 \otimes p)\mathcal{R}_0 = \mathcal{R}_0(1 \otimes p),$$

and when the dust settles, we get our formulas for ρ_1 . But QU is a quasi-triangular Hopf algebra, and hence ρ_1 is **homomorphic**. Read more at [BV1, BV2] and hear more at $\omega\epsilon\beta/\text{SolvApp}$, $\omega\epsilon\beta/\text{Dogma}$, $\omega\epsilon\beta/\text{DoPeGDO}$, $\omega\epsilon\beta/\text{FDA}$, $\omega\epsilon\beta/\text{AQDW}$.

Also, we can (and know how to) look at higher powers of ϵ and we can (and more or less know how to) replace sl_2 by arbitrary semi-simple Lie algebra (e.g., [Sch]). So ρ_1 is **not alone!**



Schaveling

These constructions are very similar to Rozansky-Overbay [Ro1, Ro2, Ro3, Ov] and hence to the "loop expansion" of the Kontsevich integral and the coloured Jones polynomial [Oh2].

If this all reads like **insanity** to you, it should (and you haven't seen half of it). Simple things should have simple explanations.

Hence, **Homework**. Explain ρ_1 with no reference to quantum voodoo and find it a topology home (large enough to house generalizations!). Make explicit the homomorphic properties of ρ_1 . Use them to do topology!

P.S. As a friend of Δ , ρ_1 gives a genus bound, sometimes better than Δ 's. How much further does this friendship extend?

A Small-Print Page on $\rho_d, d > 1$.

Definition. $\langle f(z_i), h(\zeta_i) \rangle_{\{z_i\}} := f(\partial_{\zeta_i})h|_{\zeta_i=0}$, so $\langle p^2 x^2, \mathbb{Q}^{g\pi\xi} \rangle = 2g^2$.

Baby Theorem. There exist (non unique) power series $r^\pm(p_1, p_2, x_1, x_2) = \sum_d \epsilon^d r_d^\pm(p_1, p_2, x_1, x_2) \in \mathbb{Q}[T^{\pm 1}, p_1, p_2, x_1, x_2][[\epsilon]]$ with $\deg r_d^\pm \leq 2d + 2$ (“docile”) such that the power series $Z^b = \sum \rho_d \epsilon^d :=$

$$\left\langle \exp \left(\sum_c r^s(p_i, p_j, x_i, x_j) \right), \exp \left(\sum_{\alpha\beta} g_{\alpha\beta} \pi_\alpha \xi_\beta \right) \right\rangle_{\{p_\alpha, x_\beta\}}$$

is a bnot invariant. Beyond the once-and-for-all computation of $g_{\alpha\beta}$ (a matrix inversion), Z^b is computable in $O(n^d)$ operations in the ring $\mathbb{Q}[T^{\pm 1}]$.

(Bnots are knot diagrams modulo the braid-like Reidemeister moves, but not the cyclic ones).

Theorem. There also exist docile power series $\gamma^\varphi(\bar{p}, \bar{x}) = \sum_d \epsilon^d \gamma_d^\varphi \in \mathbb{Q}[T^{\pm 1}, \bar{p}, \bar{x}][[\epsilon]]$ such that the power series $Z = \sum \rho_d \epsilon^d :=$

$$\left\langle \exp \left(\sum_c r^s(p_i, p_j, x_i, x_j) + \sum_k \gamma^{\varphi^k}(\bar{p}_k, \bar{x}_k) \right), \exp \left(\sum_{\alpha\beta} g_{\alpha\beta} (\pi_\alpha + \bar{\pi}_\alpha) (\xi_\beta + \bar{\xi}_\beta) + \sum_\alpha \pi_\alpha \bar{\xi}_\alpha \right) \right\rangle_{\{p_\alpha, \bar{p}_\alpha, x_\beta, \bar{x}_\beta\}}$$

is a knot invariant, as easily computable as Z^b .

Implementation. Data, then program (with output using the Conway variable $z = \sqrt{T} - 1/\sqrt{T}$), and then a demo. See `Rho.nb` of `omega beta alpha`.

`VerY1,ϕ[k-] = ϕ (1/2 -  $\bar{p}_k \bar{x}_k$ ); VerY2,ϕ[k-] = -ϕ2  $\bar{p}_k \bar{x}_k$  / 2;`
`VerY3,ϕ[k-] := -ϕ3  $\bar{p}_k \bar{x}_k$  / 6`

`Ver1,s[i-, j-] :=`
`s (-1+2 pi xi - 2 pj xj + (-1+Ts) pi pj xi2 + (1-Ts) pj2 xi2 - 2 pi pj xi xj + 2 pj2 xi xj) / 2`

`Ver2,1[i-, j-] :=`
`(-6 pi xi + 6 pj xi - 3 (-1+3 T) pi pj xi2 + 3 (-1+3 T) pj2 xi2 + 4 (-1+T) pi2 pj xi2 -`
`2 (-1+T) (5+T) pi pj2 xi3 + 2 (-1+T) (3+T) pj3 xi3 + 18 pi pj xi xj -`
`18 pj2 xi xj - 6 pi2 pj xi2 xj + 6 (2+T) pi pj2 xi2 xj - 6 (1+T) pj3 xi2 xj -`
`6 pi pj2 xi xj2 + 6 pj3 xi xj2) / 12`

`Ver2,-1[i-, j-] :=`
`(-6 T2 pi xi + 6 T2 pj xi + 3 (-3+T) T pi pj xi2 - 3 (-3+T) T pj2 xi2 -`
`4 (-1+T) T pi2 pj xi2 + 2 (-1+T) (1+5 T) pi pj2 xi3 - 2 (-1+T) (1+3 T) pj3 xi3 +`
`18 T2 pi pj xi xj - 18 T2 pj2 xi xj - 6 T2 pi2 pj xi2 xj + 6 T (1+2 T) pi pj2 xi2 xj -`
`6 T (1+T) pj3 xi2 xj - 6 T2 pi pj2 xi xj2 + 6 T2 pj3 xi xj2) / (12 T2)`

`Z2[GST48] (* takes a few minutes *)`

$$\begin{aligned} & \{1 - 4z^2 - 61z^4 - 207z^6 - 296z^8 - 210z^{10} - 77z^{12} - 14z^{14} - z^{16}, \\ & 1 + (38z^2 + 255z^4 + 1696z^6 + 16281z^8 + 86952z^{10} + 259994z^{12} + 487372z^{14} + 615066z^{16} + 543148z^{18} + 341714z^{20} + \\ & 153722z^{22} + 48983z^{24} + 10776z^{26} + 1554z^{28} + 132z^{30} + 5z^{32}) \epsilon + \\ & (-8 - 484z^2 + 9709z^4 + 165952z^6 + 1590491z^8 + 16256508z^{10} + 115341797z^{12} + 432685748z^{14} + 395838354z^{16} - 4017557792z^{18} - 23300064167z^{20} - \\ & 70082264972z^{22} - 142572271191z^{24} - 209475503700z^{26} - 221616295209z^{28} - 151502648428z^{30} - 23700199243z^{32} + \\ & 99462146328z^{34} + 164920463074z^{36} + 162550825432z^{38} + 119164552296z^{40} + 69153062608z^{42} + 32547596611z^{44} + 12541195448z^{46} + \\ & 3961384155z^{48} + 1021219696z^{50} + 212773106z^{52} + 35264208z^{54} + 4537548z^{56} + 436600z^{58} + 29536z^{60} + 1252z^{62} + 25z^{64}) \epsilon^2\} \end{aligned}$$

`TableForm[Table[Join[{K[[1]]K[[2]]}, Z3[K]], {K, AllKnots[{3, 6]}}], TableAlignments → Center] (* takes a few minutes *)`

3_1	$1 + z^2$	$1 + (2z^2 + z^4) \epsilon + (2 - 4z^2 + 3z^4 + 4z^6 + z^8) \epsilon^2 + (-12 + 74z^2 - 27z^4 - 20z^6 + 8z^8 + 6z^{10} + z^{12}) \epsilon^3$
4_1	$1 - z^2$	$1 + (-2 + 2z^4) \epsilon^2$
5_1	$1 + 3z^2 + z^4$	$1 + (10z^2 + 21z^4 + 12z^6 + 2z^8) \epsilon + (6 - 28z^2 + 33z^4 + 364z^6 + 655z^8 + 536z^{10} + 227z^{12} + 48z^{14} + 4z^{16}) \epsilon^2 + (-60 + 970z^2 + 645z^4 - 3380z^6 - 3280z^8 - 7470z^{10} + 19475z^{12} + 20536z^{14} + 12564z^{16} + 4774z^{18} + 1109z^{20} + 144z^{22} + 8z^{24}) \epsilon^3$
5_2	$1 + 2z^2$	$1 + (6z^2 + 5z^4) \epsilon + (4 - 20z^2 + 43z^4 + 64z^6 + 26z^8) \epsilon^2 + (-36 + 498z^2 - 883z^4 + 100z^6 + 816z^8 + 556z^{10} + 146z^{12}) \epsilon^3$
6_1	$1 - 2z^2$	$1 + (-2z^2 + z^4) \epsilon + (-4 + 4z^2 + 25z^4 - 8z^6 + 2z^8) \epsilon^2 + (12 + 154z^2 - 223z^4 - 608z^6 + 100z^8 - 52z^{10} + 10z^{12}) \epsilon^3$
6_2	$1 - z^2 - z^4$	$1 + (-2z^2 - 3z^4 + 2z^6 + z^8) \epsilon + (-2 - 4z^2 + 29z^4 + 28z^6 + 42z^8 - 8z^{10} - 2z^{12} + 4z^{14} + z^{16}) \epsilon^2 + (12 + 166z^2 + 155z^4 - 194z^6 - 2453z^8 - 1622z^{10} - 1967z^{12} - 258z^{14} + 49z^{16} - 30z^{18} + z^{20} + 6z^{22} + z^{24}) \epsilon^3$
6_3	$1 + z^2 + z^4$	$1 + (2 + 8z^2 - 16z^6 - 24z^8 - 16z^{10} - 2z^{12}) \epsilon^2$

`Ver3,1[i-, j-] :=`
`(4 pi xi - 4 pj xi + 2 (5+7 T) pi pj xi2 - 2 (5+7 T) pj2 xi2 - 4 (-5+6 T) pi2 pj xi3 +`
`4 (-16+17 T+2 T2) pi pj2 xi3 - 4 (-11+11 T+2 T2) pj3 xi3 + 3 (-1+T) pi3 pj xi4 -`
`3 (-1+T) (4+3 T) pi2 pj2 xi4 + (-1+T) (13+22 T+T2) pi pj3 xi4 -`
`(-1+T) (4+13 T+T2) pj4 xi4 - 28 pi pj xi xj + 28 pj2 xi xj + 36 pi2 pj xi2 xj -`
`12 (9+2 T) pi pj2 xi2 xj + 24 (3+T) pj3 xi2 xj - 4 pi3 pj xi3 xj + 28 T pi2 pj2 xi3 xj -`
`4 (-6+17 T+T2) pi pj3 xi3 xj + 4 (-5+10 T+T2) pj4 xi3 xj + 24 pi pj2 xi xj2 -`
`24 pj3 xi xj2 - 24 pi2 pj2 xi2 xj2 + 6 (10+T) pi pj3 xi2 xj2 - 6 (6+T) pj4 xi2 xj2 -`
`4 pi pj3 xi xj3 + 4 pj4 xi xj3) / 24`

`Ver3,-1[i-, j-] :=`
`(-4 T3 pi xi + 4 T3 pj xi - 2 T2 (7+5 T) pi pj xi2 + 2 T2 (7+5 T) pj2 xi2 -`
`4 T2 (-6+5 T) pi2 pj xi3 + 4 T (-2-17 T+16 T2) pi pj2 xi3 -`
`4 T (-2-11 T+11 T2) pj3 xi3 + 3 (-1+T) T2 pi3 pj xi4 - 3 (-1+T) T (3+4 T) pi2 pj2 xi4 +`
`(-1+T) (1+22 T+13 T2) pi pj3 xi4 - (-1+T) (1+13 T+4 T2) pj4 xi4 +`
`28 T3 pi pj xi xj - 28 T3 pj2 xi xj - 36 T3 pi2 pj xi2 xj + 12 T2 (2+9 T) pi pj2 xi2 xj -`
`24 T2 (1+3 T) pj3 xi2 xj + 4 T3 pi3 pj xi3 xj - 28 T2 pi2 pj2 xi3 xj -`
`4 T (-1-17 T+6 T2) pi pj3 xi3 xj + 4 T (-1-10 T+5 T2) pj4 xi3 xj -`
`24 T3 pi pj2 xi xj2 + 24 T3 pj3 xi xj2 + 24 T3 pi2 pj2 xi2 xj2 - 6 T2 (1+10 T) pi pj3 xi2 xj2 +`
`6 T2 (1+6 T) pj4 xi2 xj2 + 4 T3 pi pj3 xi xj3 - 4 T3 pj4 xi xj3) / (24 T3)`

`{p*, x*, p̄*, x̄*} = {π, ξ, π̄, ξ̄}; (z-)* := (z*)†;`
`Zip{}[ε-] := ε;`
`Zip{z,zs,zs...}[ε-] :=`
`(Collect[ε // Zip{zs}, z] /. f-. zd- => {D[f, {z*, d}]}]) /. z* -> 0`

`gPair[fs, w-] :=`
`gPair[fs, w] =`
`Collect[ZipJoin@Table[{pα, p̄α, xα, x̄α}, {α, w}]] [`
`(Times @@ (V /@ fs))`
`Exp[Sum[gαβ (πα + π̄α) (ξβ + ξ̄β), {α, w}, {β, w}] - Sum[ξ̄α πα, {α, w}]]],`
`g-, Factor]`

`T2z[p-] := Module[{q = Expand[p], n, c},`
`If[q == 0, 0, c = Coefficient[q, T, n = Exponent[q, T]];`
`c z2 n + T2z[q - c (T1/2 - T-1/2)2 n]]];`

`Zθ[K-] := Module[{Cs, ϕ, n, A, s, i, j, k, Δ, G, d1, Z1, Z2, Z3},`
`{Cs, ϕ} = Rot[K]; n = Length[Cs]; A = IdentityMatrix[2 n + 1];`
`Cases[Cs, {s-, i-, j-} => {A[[{i, j}, {i+1, j+1}]] += (-Ts Ts - 1)}];`
`{Δ, G} = Factor@{T(-Total[ϕ]-Total[Cs[[All,1]])/2 Det@A, Inverse@A};`
`Z1 =`
`Exp[Total[Cases[Cs, {s-, i-, j-} => Sum[ed1 rd1,s[i, j], {d1, d}]]] +`
`Sum[ed1 Yd1,ϕ[[k]][k], {k, 2 n}, {d1, d}] /. Y-,ϕ[_] -> 0];`
`Z2 = Expand[F[{}, {}] × Normal@Series[Z1, {ε, 0, d}]] /.`
`F[fs, {es---}] × (f : (r | y)ps---[is---])p- =>`
`F[Join[fs, Table[f, p]], DeleteDuplicates@{es, is}];`
`Z3 = Expand[Z2 /. F[fs, es-] => Expand[gPair[`
`Replace[fs, Thread[es -> Range@Length@es], {2}], Length@es`
`] /. gα,β => G[[es[[α], es[[β]]]]];`
`Collect[{Δ, Z3 /. ep- -> p! Δ2 p ep}, ε, T2z] ];`