



A Fast, Strong, Topologically Meaningful and Fun Knot Invariant

Abstract. The title covers all the good. The bad is that we don't really understand this invariant Θ . Wait, is that just part of the fun?

Continues Rozansky, Kricker, Garoufalidis, and Oh-tsuki [Ro1, Ro2, Ro3, Kr, GR, Oh], joint with van der Veen [BV3].

Acknowledgement. Work supported by NSERC grants RGPIN-2018-04350 and RGPIN-2025-06718 and by the Chu Family Foundation (NYC).

Strong. Testing $\Theta = (\Delta, \theta)$ on prime knots up to mirrors and reversals, counting the number of distinct values (with deficits in parenthesis):

(Vol is approximate, $H \supset \Delta, J$)

	knots	$(H, Kh, Vol, \sigma_{LT})$	$\Theta = (\Delta, \theta)$	together
xing ≤ 10	249	249 (0)	249 (0)	249 (0)
xing ≤ 11	801	787 (14)	798 (3)	798 (3)
xing ≤ 12	2,977	(84)	(19)	(10)
xing ≤ 13	12,965	(911)	(194)	(169)
xing ≤ 14	59,937	(5,917)	(1,118)	(972)
xing ≤ 15	313,230	(41,434)	(6,758)	(6,304)

Fast. Here's Θ on a random 300 crossing knot (from [DHOEBL]). For almost every other invariant, that's science fiction.


Fun. There's so much more to see in 2D pictures than in 1D ones! Yet almost nothing of the patterns you see we know how to prove. We'll have fun with that over the next few years. Would you join?

Meaningful. θ gives a genus bound (with confidence and with a near proof). θ seems to give a criterion for a knot to be fibered (conjectured with a large scale verification). There are "safe" conjectured characterizations of θ as "the two loop invariant" and as "the one cobracket invariant". We hope (with reason) θ will say something about ribbon knots.

Conventions. T, T_1 , and T_2 are indeterminates and $T_3 := T_1 T_2$.



van der Veen



Knots.

Something simple: numbers, polynomials, matrices, etc.

Tell them apart? Alternating? Bound a genus 7 surface? Complement is fibered over S^1 ? Complement is hyperbolic? Bounds a disk with only ribbon singularities? Bounds a topological / smooth non-singular disk in B^4 ? ...

Preparation. Draw an n -crossing knot K as a diagram D as on the right: all crossings face up, and the edges are marked with a running index $k \in \{1, \dots, 2n+1\}$ and with rotation numbers φ_k .

Model T Traffic Rules. Cars always drive forward. When a car crosses over a sign- s bridge it goes through with (algebraic) probability $T^s \sim 1$, but falls off with probability $1 - T^s \sim 0$. At the very end, cars fall off and disappear. On various edges traffic counters are placed. See also [Jo, LTW].



image credits: diamondtraffic.com





image credits: Dall-E

Definition. The traffic function $G = (g_{\alpha\beta})$ (also, the Green function or the two-point function) is the reading of a traffic counter at β , if car traffic is injected at α (if $\alpha = \beta$, the counter is after the injection point). There are also model- T_v traffic functions $G_v = (g_{v\alpha\beta})$ for $v = 1, 2, 3$.

Example.


$$\sum_{p \geq 0} (1-T)^p = T^{-1}$$

$$G = \begin{pmatrix} 1 & T^{-1} & 1 \\ 0 & T^{-1} & 1 \\ 0 & 0 & 1 \end{pmatrix}$$




Theorem [BV3]. With $c = (s, i, j)$, $c_0 = \begin{matrix} \uparrow s=1 \\ s_0, i_0, j_0 \end{matrix}$, $c_1 = \begin{matrix} \uparrow s=-1 \\ s_1, i_1, j_1 \end{matrix}$ denoting crossings, there is a quadratic $F_1(c) \in \mathbb{Q}(T_\nu)[g_{\nu\alpha\beta} : \begin{matrix} i & j \\ j & i \end{matrix}]$, $\alpha, \beta \in \{i, j\}$, a cubic $F_2(c_0, c_1) \in \mathbb{Q}(T_\nu)[g_{\nu\alpha\beta} : \alpha, \beta \in \{i_0, j_0, i_1, j_1\}]$, and a linear $F_3(\varphi, k)$ such that θ is a knot invariant:

$$\theta(D) := \underbrace{\Delta_1 \Delta_2 \Delta_3}_{\text{normalization, see later}} \left(\sum_c F_1(c) + \sum_{c_0, c_1} F_2(c_0, c_1) + \sum_k F_3(\varphi_k, k) \right),$$




If these pictures remind you of Feynman diagrams, it's because they are Feynman diagrams [BN2].

Lemma 1. The traffic function $g_{\alpha\beta}$ is a “relative invariant”:



Proof.



Lemma 2. With $k^+ := k + 1$, the “g-rules” hold near a crossing $c = (s, i, j)$:

$$g_{j\beta} = g_{j^+\beta} + \delta_{j\beta} \quad g_{i\beta} = T^s g_{i^+\beta} + (1 - T^s) g_{j^+\beta} + \delta_{i\beta} \quad g_{2n^+, \beta} = \delta_{2n^+, \beta}$$

$$g_{\alpha i^+} = T^s g_{\alpha i} + \delta_{\alpha i^+} \quad g_{\alpha j^+} = g_{\alpha j} + (1 - T^s) g_{\alpha i} + \delta_{\alpha j^+} \quad g_{\alpha, 1} = \delta_{\alpha, 1}$$

Corollary 1. G is easily computable, for $AG = I (= GA)$, with A the $(2n+1) \times (2n+1)$ identity matrix with additional contributions:

	A	$\text{col } i^+$	$\text{col } j^+$
$c = (s, i, j) \mapsto$			
row i	$-T^s$	$T^s - 1$	
row j	0	-1	

For the trefoil example, we have:

$$A = \begin{pmatrix} 1 & -T & 0 & 0 & T-1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -T & 0 & 0 & T-1 \\ 0 & 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & T-1 & 0 & 1 & -T & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$G = \begin{pmatrix} 1 & T & 1 & T & 1 & T & 1 \\ 0 & 1 & \frac{1}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T^2}{T^2-T+1} & 1 \\ 0 & 0 & \frac{1}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T^2}{T^2-T+1} & 1 \\ 0 & 0 & \frac{1-T}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T^2}{T^2-T+1} & 1 \\ 0 & 0 & \frac{1-T}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T}{T^2-T+1} & \frac{T^2}{T^2-T+1} & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Note. The Alexander polynomial Δ is given by

$$\Delta = T^{(-\varphi-w)/2} \det(A), \quad \text{with } \varphi = \sum_k \varphi_k, \quad w = \sum_c s.$$

We also set $\Delta_\nu := \Delta(T_\nu)$ for $\nu = 1, 2, 3$.

Questions, Conjectures, Expectations, Dreams.

Question 1. What's the relationship between Θ and the Garoufalidis-Kashaev invariants [GK, GL]?

Conjecture 2. On classical (non-virtual) knots, θ always has hexagonal (D_6) symmetry.

Conjecture 3. θ is the ϵ^1 contribution to the “solvable approximation” of the sl_3 universal invariant, obtained by running the quantization machinery on the double $\mathcal{D}(b, b, \epsilon\delta)$, where b is the Borel subalgebra of sl_3 , b is the bracket of b , and δ the cobracket. See [BV2, BN1, Sch]

Conjecture 4. θ is equal to the “two-loop contribution to the Kontsevich Integral”, as studied by Garoufalidis, Rozansky, Kricker, and in great detail by Ohtsuki [GR, Ro1, Ro2, Ro3, Kr, Oh].

Fact 5. θ has a perturbed Gaussian integral formula, with integration carried out over over a space $6E$, consisting of 6 copies of the space of edges of a knot diagram D . See [BN2].



Conjecture 6. For any knot K , its genus $g(K)$ is bounded by the T_1 -degree of θ : $2g(K) \geq \deg_{T_1} \theta(K)$.

Conjecture 7. $\theta(K)$ has another perturbed Gaussian integral formula, with integration carried out over over the space $6H_1$, consisting of 6 copies of $H_1(\Sigma)$, where Σ is a Seifert surface for K .

Expectation 8. There are many further invariants like θ , given by Green function formulas and/or Gaussian integration formulas. One or two of them may be stronger than θ and as computable.


Dream 9. These invariants can be explained by something less foreign than semisimple Lie algebras.

Dream 10. With Conjecture 7 in mind, θ will have something to say about ribbon knots.



customize.
Add fibred, HFk

Corollary 2. Proving invariance is easy:



Invariance under R3

This is Theta.nb of <http://drorbn.net/v25/ap>.

⊙ Once[<< KnotTheory` ; << Rot.m; << PolyPlot.m];

⊙ $T_3 = T_1 T_2$;

⊙ $CF[\mathcal{E}_-] := \text{Expand@Collect}[\mathcal{E}, g_-, F] /. F \rightarrow \text{Factor};$

⊙ $F_1[\{s_-, i_-, j_-\}] =$
 $CF[$
 $s (1/2 - g_{3ii} + T_2^s g_{1ii} g_{2ji} - g_{1ii} g_{2jj} -$
 $(T_2^s - 1) g_{2ji} g_{3ii} + 2 g_{2jj} g_{3ii} - (1 - T_3^s) g_{2ji} g_{3ji} -$
 $g_{2ii} g_{3jj} - T_2^s g_{2ji} g_{3jj} + g_{1ii} g_{3jj} +$
 $((T_1^s - 1) g_{1ji} (T_2^s g_{2ji} - T_2^s g_{2jj} + T_2^s g_{3jj}) +$
 $(T_3^s - 1) g_{3ji}$
 $(1 - T_2^s g_{1ii} - (T_1^s - 1) (T_2^s + 1) g_{1ji} +$
 $(T_2^s - 2) g_{2jj} + g_{2ij})) / (T_2^s - 1)]$;

⊙ $F_2[\{s0_-, i0_-, j0_-\}, \{s1_-, i1_-, j1_-\}] :=$
 $CF[s1_-(T_1^{s0} - 1) (T_2^{s1} - 1)^{-1} (T_3^{s1} - 1) g_{1,j1,i0} g_{3,j0,i1}$
 $((T_2^{s0} g_{2,i1,i0} - g_{2,i1,j0}) - (T_2^{s0} g_{2,j1,i0} - g_{2,j1,j0}))]$

⊙ $F_3[\varphi_-, k_-] = -\varphi / 2 + \varphi g_{3kk}$;

⊙ $\delta_{i_-, j_-} := \text{If}[i == j, 1, 0]$;

$gR_{s_-, i_-, j_-} := \{$
 $g_{v_j \beta_-} \Rightarrow g_{v_j^+ \beta} + \delta_{j \beta},$
 $g_{v_i \beta_-} \Rightarrow T_v^s g_{v_i^+ \beta} + (1 - T_v^s) g_{v_j^+ \beta} + \delta_{i \beta},$
 $g_{v_{-a} i^+} \Rightarrow T_v^s g_{v_{ai}} + \delta_{ai^+},$
 $g_{v_{-a} j^+} \Rightarrow g_{v_{aj}} + (1 - T_v^s) g_{v_{ai}} + \delta_{aj^+}$
 $\}$

⊙ $DSum[Cs_][c] := \text{Sum}[F_1[c], \{c, \{Cs\}\}] +$
 $\text{Sum}[F_2[c0, c1], \{c0, \{Cs\}\}, \{c1, \{Cs\}\}]$
 $lhs = DSum[\{1, j, k\}, \{1, i, k^+\}, \{1, i^+, j^+\},$
 $\{s, m, n\}] // . gR_{1,j,k} \cup gR_{1,i,k^+} \cup gR_{1,i^+,j^+};$
 $rhs = DSum[\{1, i, j\}, \{1, i^+, k\}, \{1, j^+, k^+\},$
 $\{s, m, n\}] // . gR_{1,i,j} \cup gR_{1,i^+,k} \cup gR_{1,j^+,k^+};$
 $\text{Simplify}[lhs == rhs]$

⊙ True

The Main Program

⊙ $\Theta[K_-] := \text{Module}[\{Cs, \varphi, n, A, \Delta, G, ev, \theta\},$
 $\{Cs, \varphi\} = \text{Rot}[K]; n = \text{Length}[Cs];$
 $A = \text{IdentityMatrix}[2n + 1];$
 $\text{Cases}[Cs, \{s_-, i_-, j_-\} \Rightarrow$
 $(A[\{i, j\}, \{i + 1, j + 1\}] += (-T^s T^s - 1))];$
 $\Delta = T^{(-\text{Total}[\varphi] - \text{Total}[Cs][All, 1]) / 2} \text{Det}[A];$
 $G = \text{Inverse}[A];$
 $ev[\mathcal{E}_-] :=$
 $\text{Factor}[\mathcal{E} /. g_{v_-, \alpha_-, \beta_-} \Rightarrow (G[\alpha, \beta] /. T \rightarrow T_v)];$
 $\theta = ev[\sum_{k=1}^n F_1[Cs[k]]];$
 $\theta += ev[\sum_{k1=1}^n \sum_{k2=1}^n F_2[Cs[k1], Cs[k2]]];$
 $\theta += ev[\sum_{k=1}^{2n} F_3[\varphi[k], k]];$
 Factor@
 $\{\Delta, (\Delta /. T \rightarrow T_1) (\Delta /. T \rightarrow T_2) (\Delta /. T \rightarrow T_3) \theta\}];$


The Trefoil, Conway, and Kinoshita-Terasaka

⊙ $\Theta[\text{Knot}[3, 1]] // \text{Expand}$

$$\begin{aligned} & \left\{ -1 + \frac{1}{T} + T, -\frac{1}{T_1^2} - T_1^2 - \frac{1}{T_2^2} - \frac{1}{T_1^2 T_2^2} + \frac{1}{T_1 T_2^2} + \right. \\ & \left. \frac{1}{T_1^2 T_2} + \frac{T_1}{T_2} + \frac{T_2}{T_1} + T_1^2 T_2 - T_2^2 + T_1 T_2^2 - T_1^2 T_2^2 \right\} \end{aligned}$$

⊙ $\text{GraphicsRow}[\text{PolyPlot}[\Theta[\text{Knot}[\#]]] \& /@$
 $\{"3_1", "K11n34", "K11n42"\}]$

⊙




(Note that the genus of the Conway knot appears to be bigger than the genus of Kinoshita-Terasaka)

Some Torus Knots

⊙ $\text{GraphicsRow}[\text{ImageCompose}[$
 $\text{PolyPlot}[\Theta[\text{TorusKnot}[\#]], \text{ImageSize} \rightarrow 480],$
 $\text{TubePlot}[\text{TorusKnot}[\#], \text{ImageSize} \rightarrow 240],$
 $\{\text{Right}, \text{Bottom}\}, \{\text{Right}, \text{Bottom}\}$
 $] \& /@ \{\{13, 2\}, \{17, 3\}, \{13, 5\}, \{7, 6\}\}]$



⊙



The 132-crossing torus knot $T_{22/7}$:

(many more at [ωεβ/TK](#))

Random knots from [DHOEBL] with 51 – 75 crossings: (many more at [ωεβ/DK](#))



[BN1] D. Bar-Natan, *Everything around sl_{2+}^{ϵ} is DoPeGDO. So what?*, talk in Da Nang, May 2019. Handout and video at [ωεβ/DPG](#).

[BN2] —, *Knot Invariants from Finite Dimensional Integration*, talks in Beijing (July 2024, [ωεβ/icbs24](#)) and in Geneva (August 2024, [ωεβ/ge24](#)).

[BV1] —, R. van der Veen, *A Perturbed-Alexander Invariant*, *Quantum Topology* **15** (2024) 449–472, [ωεβ/APAI](#).

[BV2] —, —, *Perturbed Gaussian Generating Functions for Universal Knot Invariants*, [arXiv:2109.02057](#).

[BV3] —, —, *A Fast, Strong, Topologically Meaningful, and Fun Knot Invariant*, [ωεβ/Theta](#) and [arXiv:2509.18456](#).

[DHOEBL] N. Dunfield, A. Hirani, M. Obeidin, A. Ehrenberg, S. Bhattacharyya, D. Lei, and others, *Random Knots: A Preliminary Report*, lecture notes at [ωεβ/DHOEBL](#). Also a data file at [ωεβ/DD](#).

[GK] ~~S. Garoufalidis, R. Kashaev, *Multivariable Knot Polynomials from Braided Hopf Algebras with Automorphisms*, [arXiv:2311.11528](#).~~

[GL] —, S. Y. Li, *Patterns of the V_2 -polynomial of knots*, [arXiv:2409.03557](#).

[GR] —, L. Rozansky, *The Loop Expansion of the Kontsevich Integral, the Null-Move, and S-Equivalence*, [arXiv:math.GT/0003187](#).

[Jo] V. F. R. Jones, *Hecke Algebra Representations of Braid Groups and Link Polynomials*, *Annals Math.*, **126** (1987) 335–388.

[Kr] A. Kricker, *The Lines of the Kontsevich Integral and Rozansky’s Rationality Conjecture*, [arXiv:math/0005284](#).

[LTW] X-S. Lin, F. Tian, Z. Wang, *Bureau Representation and Random Walk on String Links*, *Pac. J. Math.*, **182-2** (1998) 289–302, [arXiv:q-alg/9605023](#).

[Oh] T. Ohtsuki, *On the 2-loop Polynomial of Knots*, *Geom. Top.* **11** (2007) 1357–1475.

[Ov] A. Overbay, *Perturbative Expansion of the Colored Jones Polynomial*, Ph.D. thesis, University of North Carolina, Aug. 2013, [ωεβ/Ov](#).

[Ro1] L. Rozansky, *A Contribution of the Trivial Flat Connection to the Jones Polynomial and Witten’s Invariant of 3D Manifolds, I*, *Comm. Math. Phys.* **175-2** (1996) 275–296, [arXiv:hep-th/9401061](#).

[Ro2] —, *The Universal R-Matrix, Bureau Representation and the Melvin-Morton Expansion of the Colored Jones Polynomial*, *Adv. Math.* **134-1** (1998) 1–31, [arXiv:q-alg/9604005](#).

[Ro3] —, *A Universal $U(1)$ -RCC Invariant of Links and Rationality Conjecture*, [arXiv:math/0201139](#).

[Sch] S. Schaveling, *Expansions of Quantum Group Invariants*, Ph.D. thesis, Universiteit Leiden, September 2020, [ωεβ/Scha](#).